

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
26 May 2005 (26.05.2005)


PCT

(10) International Publication Number
WO 2005/046583 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number:
PCT/US2004/035536
- (22) International Filing Date: 25 October 2004 (25.10.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/516,647 30 October 2003 (30.10.2003) US
- (71) Applicant (for all designated States except US): **EMORY UNIVERSITY** [US/US]; 1784 North Decatur Road, Atlanta, GA 30322 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **LOLLAR, John, S.** [US/US]; 2568 Oak Crossing Drive, Decatur, GA 30033 (US).
- (74) Agents: **CURTIS, Michael, J.** et al.; Greenlee, Winner and Sullivan, P.C., 4875 Pearl East Circle, Suite 200, Boulder, CO 80301 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: MODIFIED FVIII HAVING REDUCED IMMUNOGENICITY THROUGH MUTAGENESIS OF A2 AND C2 EPI-TOPIES



(57) Abstract: Specific amino acid loci of human fVIII interact with inhibitory antibodies of hemophilia patients after being treated with fVIII. Modified fVIII is disclosed in which the amino acid sequence is changed by multiple substitutions in human fVIII A2 and C2 domains. The modified fVIII is useful for hemophiliacs, either to avoid or prevent the action of inhibitory antibodies.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Modified fVIII Having Reduced Immunogenicity Through Mutagenesis of A2 and C2 Epitopes

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 60/516,647, filed October 30, 2003, which is incorporated herein to the extent that there is no inconsistency with the present disclosure.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] Not Applicable

BACKGROUND OF THE INVENTION

[0003] This invention relates generally to a modified mammalian factor VIII ("fVIII" herein) having amino acid substitutions which reduce its immunogenicity and/or antigenicity as compared to the proteins from which they were derived or other fVIII preparations such as human fVIII.

[0004] Blood clotting begins when platelets adhere to the cut wall of an injured blood vessel at a lesion site. Subsequently, in a cascade of enzymatically regulated reactions, soluble fibrinogen molecules are converted by the enzyme thrombin to insoluble strands of fibrin that hold the platelets together in a thrombus. At each step in the cascade, a protein precursor is converted by a protease that cleaves the next protein precursor in the series. Co-factors are required at most of the steps.

[0005] FVIII circulates as an inactive precursor in blood, bound tightly and non-covalently to von Willebrand factor. FVIII is proteolytically activated by thrombin and by factor Xa. Activation dissociates fVIII from von Willebrand factor and activates its procoagulant function in the cascade. In its active form, the protein fVIIIa is a cofactor that increases the catalytic efficiency of factor IXa toward factor X activation by several orders of magnitude.

[0006] People with deficiencies in fVIII (hemophilia A) or antibodies against fVIII who are not treated with fVIII suffer uncontrolled internal bleeding that may cause a range of serious symptoms, from inflammatory reactions in joints to early death. Severe hemophiliacs, who number about 10,000 in the United States, can be treated with infusion of human fVIII, which will restore the blood's normal clotting ability if administered with sufficient frequency and concentration. The functional definition of fVIII is that substance present in normal blood plasma that corrects the clotting defect in plasma derived from individuals with hemophilia A. Human fVIII has been characterized as a 2332 amino acid polypeptide, the amino acid sequence of which is given in SEQ ID NO: 2, having structural domains labeled as A1-A2-B-A3-C1-C2 (Vehar, G.A. et al., 1984, *Nature*, 312:337-342).

[0007] Although SEQ ID NO: 2 is the amino acid sequence of a human fVIII deduced from DNA sequencing from a human source, the term "human fVIII" is used herein to include a variety of sequence variations. For example, allelic variations in the amino-acid sequences of naturally-occurring proteins are well known and may be characterizing features of different populations or ethnic groups. Single nucleotide polymorphisms (SNP's) among individuals are well-recognized phenomena, some of which can result in coding changes and resulting variations in amino-acid sequence. In addition, it is known that many parts of the canonical fVIII sequence (SEQ ID NO: 2) can be deliberately changed by amino acid substitutions, by deleting segments of the canonical sequence, or by insertion into the canonical sequence. All or part of the B domain, from amino acids 741 to 1648, can be deleted without apparent loss of normal function. (Toole et al. (1986) *Proc. Natl. Acad. Sci. USA* 83(16):5939-5942; Eaton et al. (1986) *Biochemistry* 25(26):8343-8347; Langer et al. (1988) *Behring Inst. Mitt.* 82:16-25; Meulien et al. (1988) *Protein Eng.* 2(4):301-6; and U.S. patent 4,868,112, all of which are incorporated herein by reference.) Specific amino-acid substitutions can be made at many sites without significant loss of procoagulant activity. (U.S. patents 5,744,446; 5,859,204; 6,060,447; 6,180,371; 6,228,620; and 6,376,463.) Furthermore, comparisons between the amino acid sequences of human and animal fVIII's demonstrate a high degree of conservation of sequence in domains other than the B domain. (Diamond et al. (1992) *Hum Mutat* 1(3):248-57; Elder et al. (1993) *Genomics* 16(2):374-9; Healy et al. (1996) *Blood* 88(11):4209-14; Cameron et al. (1998) *Thromb Haemost* 79(2):317-22; Watzka et al. (2004) *Thromb*

Haemost 91(1):38-42; and U.S. patents 5,364,771 and 5,859,204) Therefore, persons skilled in the art can reasonably predict where amino-acid substitutions are most likely to be tolerated in terms of coagulant activity, where substitutions of like-for-like amino acids can be made while retaining coagulant activity and where substitutions may result in loss of coagulant activity.

[0008] The term "human fVIII" as used herein, includes all such variations from the canonical sequence of SEQ ID NO: 2 as may exist, be discovered or artificially induced, provided the essential functional attribute of procoagulant activity in a human Hemophillia A patient, sufficient for therapeutic efficacy, is maintained. Procoagulant activity is commonly expressed in units/ml of plasma, measured by either a coagulation assay or a chromogenic assay comparing a sample of a patient's plasma with an international reference standard. The activity of normal plasma is about 1 unit/ml. A "human fVIII" as defined herein will have procoagulant activity to provide at least 0.01 units/ml plasma 15 minutes after administration to a hemophiliac patient lacking inhibitor antibodies. Although the number of changes from the canonical sequence is likely to be small, human fVIII, as herein defined, can be distinguished from the other animal fVIII's by simple sequence identity comparison. Thus, a given sequence of human fVIII will have more amino acids identical to canonical SEQ ID NO: 2 than to any known animal sequence, after taking into account corresponding deletions, e.g. in the B domain.

[0009] The development of antibodies ("inhibitors" or "inhibitory antibodies") that inhibit the activity of fVIII is a serious complication in the management of patients with hemophilia. Inhibitory antibodies (inhibitors) to fVIII either develop as alloantibodies in hemophilia A patients following fVIII infusions or as autoantibodies in nonhemophiliacs (Hoyer, L.W. and D. Scandella, 1994, *Semin.Hematol.* 31:1-5). Antibodies develop in approximately 20% of patients with hemophilia A in response to therapeutic infusions of fVIII. In previously untreated patients with hemophilia A who develop inhibitors, the inhibitors usually develop within one year of treatment. Additionally, autoantibodies that inactivate fVIII occasionally develop in individuals with previously normal fVIII levels. Antibodies to epitopes in the A2, *ap*-A3, and C2 domains within the A1-A2-B-*ap*-A3-C1-C2 fVIII molecule are responsible for all

anticoagulant activity in most inhibitor plasmas (Prescott, R. et al., 1997, *Blood* 89:3663-3671; Barrow, R.T. et al., 2000, *Blood* 95:557-561).

[0010] Anti-A2 inhibitors inhibit the function of activated fVIII within the intrinsic pathway factor X activation complex, apparently by blocking the binding of factor X to the complex (Lollar, P. et al., 1994, *J. Clin. Invest.* 93:2497-2504). The A2 epitope has been localized to a single, continuous sequence bounded by residues R484-1508 (Healey, J.F. et al., 1995, *J. Biol. Chem.* 270:14505-14509).

[0011] The 18-kDa C2 domain, defined as residues Ser2173 - Tyr2332 in single chain human fVIII, contains a phospholipid membrane-binding site that is necessary for the normal procoagulant function of fVIII. Human C2-specific anti-fVIII antibodies inhibit this interaction (Arai, M. et al., 1989, *J. Clin. Invest.* 83:1978-1984). Consistent with this, binding to phospholipids protects fVIII from inactivation by fVIII inhibitors (Arai et al., *supra*; Barrowcliffe, T.W. et al., 1983, *J. Lab. Clin. Med.* 101:34-43). The C2 domain also contains part of the von Willebrand factor (vWf) binding site (Saenko, E.L. et al., 1994, *J. Biol. Chem.* 269:11601-11605; Saenko, E.L. and Scandella, D., 1997, *J. Biol. Chem.* 272:18007-18014). Some inhibitors may act by interfering with this interaction (Shima, M. et al., 1995, *Br. J. Haematol.* 91:714-721; Saenko, E.L. et al., 1996 *J. Biol. Chem.* 271:27424-27431; Gilles, J.G. et al., 1999, *Thromb. Haemost.* 82:40-45).

[0012] Patients who develop antibodies to human fVIII can be managed by increasing the dose of fVIII provided the inhibitor titer is low enough. However, often the inhibitor titer is so high that it cannot be overwhelmed by increased amounts of fVIII. An alternative strategy is to bypass the need for fVIII during normal hemostasis using factor IX complex preparations (for example, KONYNE[®], Proplex[®]) or using recombinant human fVIIIa. Additionally, since porcine fVIII usually has substantially less reactivity with inhibitors than human fVIII, a partially purified porcine fVIII preparation (HYATE:C[®]) is used. Many patients who have developed inhibitory antibodies to human fVIII have been successfully treated with porcine fVIII and have tolerated such treatment for long periods of time. However, administration of porcine fVIII is not a complete solution because inhibitors occasionally develop to porcine fVIII after one or more infusions.

[0013] Hybrid fVIII molecules which substitute regions of human fVIII with the corresponding regions from animals are well known in the art. U.S. patent 5,888,974 (Lollar et al.) discloses hybrid procoagulant fVIII produced by the isolation and recombination of human with non-human fVIII subunits or domains. Similarly, U.S. patents 5,663,060 and 5,583,209 describe hybrid fVIII comprising combinations of non-human and human heavy chain and light chain subunits. U.S. patent 5,364,771 describes purified hybrid fVIII comprised of human and porcine combinations of the heavy and light subunits, including a human fVIII with a porcine A2 domain substituted for the human A2 domain.

[0014] Several preparations of human plasma-derived fVIII of varying degrees of purity are available commercially for the treatment of hemophilia A. These include a partially-purified fVIII derived from the pooled blood of many donors that is heat- and detergent-treated to inactivate viruses but contain a significant level of antigenic proteins; and a monoclonal antibody-purified fVIII that has lower levels of antigenic impurities and viral contamination. Another alternative product is recombinant human fVIII (currently sold under the trade name Refacto®) which would be free of viral contaminants. Unfortunately, human fVIII is unstable at physiologic concentrations and pH, is present in blood at an extremely low concentration (0.2 µg/ml plasma), and has low specific clotting activity. A significant proportion of patients receiving recombinant human fVIII develop inhibitory antibodies to that product.

[0015] Hemophiliacs require daily replacement of fVIII to prevent bleeding and the resulting deforming hemophilic arthropathy. However, supplies have been inadequate and problems in therapeutic use occur due to difficulty in isolation and purification, immunoreactivity, and the necessity of removing the AIDS and hepatitis infectivity risk. The use of recombinant human fVIII, partially-purified porcine fVIII, or human-animal hybrid fVIII will not resolve all the problems.

[0016] The problems associated with the commonly used, commercially available, plasma-derived fVIII have stimulated significant interest in the development of a better fVIII product. There is a need for fVIII product having greater specific activity so that more units of clotting activity can be delivered per milligram of total protein; a fVIII molecule that is stable at a selected pH and

physiologic concentration; a fVIII molecule that is less apt to cause production of inhibitory antibodies; and a fVIII molecule that evades immune detection in patients who have already acquired antibodies to human fVIII.

[0017] Reduction of antigenicity (inhibition by reaction with antibodies) is described in U.S. patents 6,180,371 and 5,744,446 which describe modified fVIII having amino acid substitutions in the A2 domain. U.S. patents 6,376,463 and 5,859,204 disclose site specific replacement of amino acids in the 484-509 region in the A2 domain of human fVIII. These patents do not disclose or suggest specific amino acid substitutions in the C2 domain which reduce antigenicity or immunogenicity as compared to wild-type fVIII or the corresponding recombinant fVIII. These patents also do not disclose the specific triple mutant in the A2 domain described in the present invention.

[0018] U.S. patent 6,770,744 discloses modified fVIII with site specific amino acid substitutions in the C2 domain at positions 2199, 2200, 2223, 2227, 2251, 2252 relative to human fVIII. The substitutions are disclosed to reduce antigenicity with respect to certain antibodies directed to the C2 domain. The 6,770,744 patent discloses a quadruple mutant having amino acid substitutions at each of positions 2199, 2200, 2223 and 2227, but does not disclose the specific quadruple mutant described in the present invention.

[0019] Pratt et al. (1999, *Nature* 402:439-442) have reported an X-ray crystallography structure of the C2 domain of human fVIII at 1.5A resolution. Pratt et al. reported that the structure partly explains why mutations in the C2 region of fVIII lead to bleeding disorders. In fact, 21 residues in the C2 region were reported to be sites of deleterious point mutations in patients with hemophilia A. Shima et al. report C2 directed inhibitory antibodies that interfere with fVIII with respect to phospholipid and Von Willebrand factor binding. Thus, it is taught by Pratt et al. that C2 inhibitors, i.e., those related to some bleeding disorders in individuals with hemophilia A, interfere with the binding of the C2 domain to phospholipid and Von Willebrand factor. This conclusion, combined with their determination that M2199, F2200, V2223, K2227, L2251 and L2252 appear at the protein-phospholipid interface, suggests that these amino acids are important for normal fVIII activity and mutation of these residues would lead to detrimentally altered phospholipid and/or Von

Willebrand binding along with an associated increase in bleeding disorders. However, U.S. patent 6,770,744 discloses amino acid substitutions at amino acid positions M2199, F2200, V2223, K2227, L2251 and L2252 which result in reduced antigenicity while maintaining coagulant activity.

[0020] U.S. patents 6,180,371; 5,888,974; 5,859,204; 5,744,446; 5,663,060; 5,583,209; and 5,364,771 and U.S. patent application 10/131,510 (all of which are incorporated herein by reference) do not disclose the specific triple mutant in the A2 domain described in the present invention. Nor do they disclose amino acid substitutions in the C2 domain of fVIII which reduce the immunogenicity as compared to wild-type fVIII or the corresponding recombinant fVIII. U.S. patent 6,770,744 (also incorporated herein by reference) discloses other amino acid substitutions in the C2 domain, but does not disclose specific amino acid substitutions in the A2 domain for reducing immunogenicity, nor the specific quadruple mutant in the C2 domain described in the present invention. None of the above patents or applications discloses a mutant combining site specific amino acid substitutions in both the A2 and C2 domain.

[0021] It is not clear from previous studies which amino acid residues and corresponding substitutions would lead to improved fVIII molecules. As described further below, one modified fVIII that contains the R484A/R489A mutation has no significant difference in immunogenicity compared to normal human fVIII, while a different fVIII that contains the R484A/R489A/P492A mutant has significantly lower immunogenicity than normal human fVIII. Similarly, three specific fVIII mutants described below, designated A2C2epi1, A2C2epi2 and A2C2epi3, have the R484A/R489A/P492A mutant in the A2 domain combined with additional multiple amino acid substitutions in C2 domain. The amino acid substitutions in the C2 domain differ between A2C2epi1, A2C2epi2, and A2C2epi3 by no more than two amino acids. Despite the fact that all three of these fVIII mutants contain the R484A/R489A/P492A mutant, which has lower immunogenicity than normal human fVIII, one fVIII mutant, A2C2epi2, does not have reduced immunogenicity compared to normal human fVIII. The other two mutants have lower immunogenicity than normal human fVIII, but only the A2C2epi3 fVIII mutant has significantly lower immunogenicity than the fVIII mutant having only the R484A/R489A/P492A mutant.

[0022] It is an object of the present invention to provide a fVIII that corrects hemophilia in a patient deficient in fVIII and can be administered routinely with a reduced risk of developing inhibitory antibodies to fVIII compared to currently available products. It is a further object of the present invention to provide methods for treatment of hemophiliacs. It is still another object of the present invention to provide a fVIII that is stable at a selected pH and physiologic concentration. It is yet another object of the present invention to provide a fVIII that has greater or more prolonged specific coagulant activity than human fVIII.

SUMMARY OF THE INVENTION

[0023] The present invention generally relates to compositions comprising recombinant human fVIII. The compositions of the invention comprise isolated, purified recombinant human fVIII molecules with coagulant activity wherein the recombinant fVIII has amino acid substitutions in the A2 and C2 domains which reduce immunogenicity and optionally antigenicity as compared to the proteins from which they were derived or other fVIII preparations. DNA sequences encoding the novel compositions of the invention as well as methods of producing the novel compositions comprising fVIII are also provided. Methods of treating patients in need of treatment with fVIII are also within the scope of this invention.

[0024] A first embodiment of the invention provides a modified fVIII comprising the C2 domain of human fVIII with a quadruple amino acid substitution in the C2 domain. The amino acid substitutions in the C2 domain of the modified recombinant fVIII reduce its capacity to elicit inhibitory antibodies as compared to the proteins from which they were derived or other available fVIII preparations. The novel composition of this embodiment is a modified fVIII molecule having amino acid substitutions in the C2 domain at each of positions 2199, 2200, 2251 and 2252.

[0025] A further embodiment is a modified fVIII having the quadruple mutant M2199L/F2200L/L2251V/ L2252F (leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252). M2199L/F2200L/L2251V/ L2252F is referenced to the human fVIII numbering system wherein amino acid

number 1 is the amino terminal alanine of mature fVIII. Amino acids will be identified herein using the generally accepted single letter code.

[0026] In one embodiment, the fVIII having the quadruple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 2173-2332. In another embodiment, the fVIII having the quadruple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1649-2332. In another embodiment, the fVIII having the quadruple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-740 and 1649-2332. In another embodiment, the fVIII having the quadruple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-745 and 1640-2332. In another embodiment, the fVIII having the quadruple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-2332. The modified fVIII has reduced immunogenicity and antigenicity to an inhibitory antibody as compared to unmodified fVIII and may lack part or the entire B domain. The modified fVIII comprises an A1/A2/A3-C1-C2 heterotrimer, an A1-A2/A3-C1-C2 heterodimer, or a single continuous polypeptide. This invention also provides pharmacological compositions comprising the above modified fVIII. This invention also provides DNA encoding the above modified fVIII and methods of making the modified fVIII by expressing said DNA. The corresponding nucleotide sequence encoding human fVIII is disclosed in SEQ ID NO: 1.

[0027] One embodiment of the present invention is a polypeptide comprising an amino acid sequence having the quadruple mutant M2199L/F2200L/L2251V/L2252F and having at least about 85% sequence homology, more usually at least about 95% sequence homology, with the C2 domain (amino acids 2173-2332) set forth in SEQ ID NO: 2.

[0028] U.S. patent 6,770,744 discloses substitutions at positions homologous to human fVIII including, but not limited to, M2199, F2200, V2223, K2227, L2251, and L2252. Of particular note, U.S. patent 6,770,744 discloses substituting isoleucine for M2199. However, a modified fVIII having the quadruple mutant M2199I/F2200L/L2251V/L2252F, as disclosed in the 6,770,744 patent, did not have significantly lower immunogenicity compared to normal human fVIII. In contrast, a modified fVIII having the novel composition of this embodiment, the quadruple mutant

M2199L/F2200L/L2251V/L2252F, which substitutes leucine for M2199 instead of isoleucine, does have significantly reduced immunogenicity compared to normal human fVIII and retains coagulant activity.

[0029] Another embodiment of this invention provides a modified fVIII comprising the A2 domain of human fVIII with a triple amino acid substitution in the A2. The amino acid substitutions in the A2 domain of the modified fVIII reduce its immunogenicity compared to the proteins from which they were derived or other available fVIII preparations. The novel composition of this embodiment is a modified fVIII having immunoreactivity reducing amino acid substituted in the A2 domain at each of positions 484, 489 and 492.

[0030] A further embodiment is a modified fVIII having the triple mutant R484A/R489A/ P492A, where alanine is substituted for arginine 484, arginine 489, and proline 492. R484A/ R489A/P492A is referenced to the human fVIII numbering system wherein amino acid number 1 is the amino terminal alanine of mature fVIII.

[0031] A further embodiment is a modified fVIII having the corresponding porcine amino acids substituted for arginine 484, arginine 489, and proline 492. The amino acid and cDNA sequences of porcine fVIII are set forth in SEQ ID NO: 3 and SEQ ID NO: 4, respectively. In SEQ ID NO: 4, the coding region begins at nucleotide position 195, the triplet GCC being the codon for amino acid number 1 (Ala) of the mature protein as given in SEQ ID NO: 3.

[0032] In one embodiment, the fVIII having the triple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 373-740. In another embodiment, the fVIII having the triple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-740. In another embodiment, the fVIII having the triple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-740 and 1649-2332. In another embodiment, the fVIII having the triple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-745 and 1640-2332. In another embodiment, the fVIII having the triple mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-2332. The modified fVIII has reduced immunogenicity and antigenicity to an inhibitory antibody as compared to unmodified fVIII and may lack

part or the entire B domain. The modified fVIII comprises an A1/A2/A3-C1-C2 heterotrimer, an A1-A2/A3-C1-C2 heterodimer, or a single continuous polypeptide. This invention also provides pharmacological compositions comprising the above modified fVIII. This invention also provides DNA encoding the above modified fVIII and methods of making the modified fVIII by expressing said DNA. The corresponding nucleotide sequence encoding human fVIII is disclosed in SEQ ID NO 1.

[0033] One embodiment of the present invention is a polypeptide comprising an amino acid sequence having the triple mutant R484A/R489A/P492A and having at least about 85% sequence homology, more usually at least about 95% sequence homology, with the A2 domain (amino acids 373-740) set forth in SEQ ID NO: 2.

[0034] Another embodiment of this invention provides a modified fVIII comprising the A2 and C2 domains of human fVIII with a triple amino acid substitution in the A2 domain combined with the previously described quadruple amino acid substitution in the C2 domain. The amino acid substitutions in both the A2 and C2 domains of the modified fVIII reduce immunogenicity of the modified fVIII when compared to the proteins from which they were derived or other available fVIII preparations having amino acid substitutions in either the A2 domain or C2 domain. The novel composition of this embodiment is modified fVIII molecule having amino acid substitutions in the C2 domain at each of positions 2199, 2200, 2251 and 2252, and having amino acid substitutions in the A2 domain at each of positions 484, 489 and 492.

[0035] A further embodiment of the present invention is a modified human fVIII molecule having the mutation R484A/R489A/P492A/M2199L/F2200L/L2251V/L2252F, where alanine is substituted for arginine 484, arginine 489, and proline 492, leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252.

[0036] In one embodiment, the fVIII having the R484A/R489A/P492A/M2199L/F2200L/L2251V/L2252F mutation comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 373-740 and 2173-2332. In another embodiment,

the fVIII having the mutation comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-740 and 1649-2332. In another embodiment, the fVIII having the mutant comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-745 and 1640-2332. In another embodiment, the fVIII having the mutation comprises the amino acid sequence set forth in SEQ ID NO: 2 from amino acids 1-2332. The modified fVIII has reduced immunogenicity and antigenicity to an inhibitory antibody as compared to unmodified fVIII and may lack part or the entire B domain. The modified fVIII comprises an A1/A2/A3-C1-C2 heterotrimer, an A1-A2/A3-C1-C2 heterodimer, or a single continuous polypeptide. This invention also provides pharmacological compositions comprising the above modified fVIII. This invention also provides DNA encoding the above modified fVIII and methods of making the modified fVIII by expressing said DNA. The corresponding nucleotide sequence encoding human fVIII is disclosed in SEQ ID NO: 1.

[0037] One embodiment of the present invention is a polypeptide comprising an amino acid sequence having the mutant R484A/R489A/P492A/M2199L/F2200L/L2251V/L2252F and having at least about 85% sequence homology, more usually at least about 95% sequence homology, with the A2 domain (amino acids 373-740) and C2 domain (amino acids 2173-2332) set forth in SEQ ID NO: 2.

[0038] In one embodiment of the present invention, the modified fVIII comprises the A1, A2, A3, C1 and C2 domains of human fVIII. In another embodiment of the present invention, the modified fVIII comprises the C2 domain of porcine fVIII and the A2 domain of human fVIII having the above mentioned amino acid substitutions at arginine 484, arginine 489, and proline 492. In another embodiment, the modified fVIII comprises the A2 domain of porcine fVIII and the C2 domain of human fVIII having the previously described quadruple amino acid substitution in the C2 domain. The amino acid sequence for porcine fVIII is set forth in SEQ ID NO: 3.

[0039] Another embodiment of the invention provides DNA sequences comprising coding sequences for the modified fVIII of the invention. Yet another embodiment of the invention provides methods of producing the modified fVIII of the invention.

[0040] fVIII of the present invention may exist as a heterodimer or heterotrimer. As a result, the separate domains and subunits do not necessarily have to be

expressed from the same DNA segment. For example, U.S. patents 6,060,447 and 6,228,620 disclose separately expressing a fVIII heavy chain (amino acids 1-740) and a fVIII light chain (amino acids 1649-2332). One embodiment of the present invention is a DNA complex comprising: (a) a first DNA segment encoding the A2 domain or A1 and A2 domains where alanine is substituted for arginine 484, arginine 489, and proline 492; and (b) a second DNA segment encoding the C2 domain, or A3, C1, and C2 domains where leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252.

[0041] The invention provides a method for reducing the immunogenicity of fVIII as well as a recombinant fVIII having reduced immunogenicity produced by the method. In particular, modified human fVIII, and methods of making such molecules, having immunoreactivity reducing amino acid substitutions in the A2 and C2 domain are described. "Immunoreactivity reducing" amino acids are defined herein as those amino acids which do not significantly contribute to an antigen-antibody interaction. Non-limiting examples of some amino acids known to be immunoreactivity-reducing include alanine, methionine, leucine, serine, and glycine. It will be understood that the reduction of immunoreactivity achievable by a given amino acid substitution will also depend on any effects the substitution may have on protein conformation, epitope accessibility and the like.

[0042] Also provided are pharmaceutical compositions and methods for treating patients having fVIII deficiency comprising administering modified recombinant fVIII.

BRIEF DESCRIPTION OF THE DRAWINGS

[0043] Figure 1 shows diagrams of various FVIII constructs and their structural domains. Human, porcine and murine forms of B domain deleted fVIII are designated HBD, PBD, and MBD, respectively. Bold type in the amino acid alignments corresponds to amino acid mutagenesis or substitution within the 484-508 A2 epitope that differs from human fVIII. Included are amino acid sequences of the A2 epitope for human B domain deleted constructs having a double amino acid substitution or a triple amino acid substitution. Also shown is a human construct designated HP9 where the 484-508 A2 epitope is substituted with the corresponding

porcine amino acid sequence. Also shown is a murine construct designated HM1 where the 484-508 A2 epitope is substituted with the corresponding human amino acid sequence.

[0044] Figure 2 shows the antigenicity of the human R484-I508 A2 segments in human B domain deleted (HBD) immunized hemophilia A mice. Plasmas were obtained from hemophilia A mice immunized with HBD and were assayed for inhibitory anti-fVIII antibodies against HBD and HP9 by Bethesda assay (A) and anti-fVIII antibodies against MBD and HM1 by ELISA (B) as described in Example 1. Each data pair corresponds to individual mouse plasma from an HBD-immunized mouse.

[0045] Figure 3 shows the correlation between the reduced antigenicity of HP9 and the increased antigenicity of HM1 in HBD-immunized hemophilia A mice. Ratios of assay titers for HBD to HP9 and for HM1 to MBD from the experiment shown in Fig. 2 were calculated and plotted. The regression line corresponds to a coefficient of correlation, r , of 0.61, which is significantly greater than zero ($p=0.001$, t test).

[0046] Figure 4 shows the comparative immunogenicity of HBD, R484A/R489A/P492A and R484A/R489A. Hemophilia A mice received intravenous injections of HBD, the triple mutant R484A/R489A/P492A or the double mutant R484A/R489A and then were tested for inhibitory anti-fVIII antibodies by Bethesda assay as described in Materials and Methods. The horizontal lines represent the sample means. The statistical parameters of the samples (mean \pm s.d.) were 670 ± 500 , 320 ± 310 , and 780 ± 570 for the HBD, R484A/R489A/P492A, and R484A/R489A groups, respectively.

[0047] Figure 5 shows in vivo clearance of R484A/R489A/P492A and HBD. Mice were injected with 100 U/kg R484A/R489A/P492A (open circles) or HBD (closed circles) by tail vein and assayed for fVIII activity by chromogenic assay described in Materials and Methods. A different group of three mice was used at each time point. Data represent the mean and standard deviation.

[0048] Figure 6 shows candidate low immunogenicity fVIII C2 domain mutations. The amino acid substitutions at phospholipid binding loops 1 and 3 of the C2 epi1, C2 epi2 and C2 epi3 mutants are shown compared to the native human amino acids.

[0049] Figure 7 shows the expression of fVIII constructs from BHK-derived cells. FVIII coagulant activity from triple-flask supernatants was assayed at the indicated times during the production runs. In this figure C2epi1, C2epi2 and C2epi3 refer to A2C2epi1, A2C2epi2 and A2C2epi3, respectively.

[0050] Figure 8 shows the SDS-PAGE of fVIII constructs. Purified HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 (1.3 µg), with and without treatment thrombin (IIa), underwent 4-15% gradient SDS-PAGE, followed by Gel-Code Blue staining. SC, single chain; H, heavy chain (A1-A2); LC, light chain (*ap*-A3-C1-C2); LC_{IIa}, thrombin-cleaved light chain (A3-C1-C2); *, low molecular weight contaminant. In this figure C2epi1, C2epi2 and C2epi3 refer to A2C2epi1, A2C2epi2 and A2C2epi3, respectively.

[0051] Figure 9 shows the Bethesda assay of the comparative immunogenicity of HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 in hemophilia mice. Mice received intravenous injections of fVIII preparations and then were tested for inhibitory anti-fVIII antibodies by Bethesda assay as described in Example 2. The horizontal lines represent the sample means.

[0052] Figure 10 shows the ELISA analysis of comparative immunogenicity of HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3. Mice received intravenous injections of fVIII preparation and then were tested by Elisa as described in Example 2. The horizontal lines represent the sample means.

[0053] Figure 11 shows the single human domain hybrid human/porcine fVIII constructs. The shaded areas designate domains having amino acid sequences corresponding to porcine fVIII. The white areas designate domains having amino acid sequences corresponding to human fVIII.

[0054] Figure 12 shows the domain specific ELISA titers from mice immunized with HBD, A2epi7 or A2C2epi3. Plasmas from ten mice selected from each of the HBD and A2epi7 groups and all five ELISA positive mice in the A2C2epi3 group

were subjected to ELISA on single human domain hybrid human/porcine fVIII coated plates as described in Example 2.

[0055] Figure 13 shows the in vitro recovery of HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3. Human hemophilia A plasma was reconstituted with purified HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 followed by assay of fVIII coagulant as described in Example 2. Data are expressed as means and sample standard deviations of expected recovery based on the nominal concentrations of purified proteins and result from at least eight samples done on separate days.

[0056] Figure 14 shows the concentration dependence of apt-reagent on the clotting times of human hemophilia A plasma reconstituted with HBD and A2C2epi3. Human hemophilia A plasma was reconstituted with HBD or A2C2epi3 and clotting times were determined as in standard one-stage fVIII assays, except the concentration of a PTT-reagent was varied. The data represents means and ranges of two independent experiments in which duplicate samples were assayed.

DETAILED DESCRIPTION OF THE INVENTION

[0057] The present invention generally relates to compositions comprising recombinant human fVIII. The compositions of the present invention comprise isolated, purified recombinant human fVIII molecules with coagulant activity. It was previously discovered that mutations in the C2 domain of fVIII reduced the binding of inhibitory antibodies of the mutants as compared to the proteins from which they were derived and/or other fVIII preparations. Novel compositions of the invention comprise recombinant human fVIII with specific amino acid substitutions in the C2 domain which reduce immunogenicity or antigenicity as compared to the proteins from which they were derived or other available fVIII preparations. Furthermore, it has been previously discovered that mutations in the A2 domain of fVIII reduce the binding of inhibitory antibodies of the mutants as compared to the proteins from which they were derived and/or other fVIII preparations. Novel compositions of the invention comprise recombinant factor human VIII with specific amino acid substitutions in the A2 domain which reduce immunogenicity or antigenicity as compared to the proteins from which they were derived or other available fVIII preparations. Novel compositions of the invention also comprise recombinant

human fVIII with specific amino acid substitutions in both the A2 and C2 domains which reduce immunogenicity or antigenicity as compared to the proteins from which they were derived or other available fVIII preparations.

[0058] Related embodiments of the invention provide for methods of treating patients in need of fVIII treatment, methods of producing the novel recombinant fVIII compositions of the invention, DNA sequences comprising coding sequences of the novel recombinant fVIII proteins, and pharmaceutical compositions comprising the novel fVIII proteins.

[0059] The present invention further provides active recombinant hybrid fVIII molecules or fragments thereof, the nucleic acid sequences encoding these hybrids, methods of preparing and isolating them, and methods for characterizing them. These hybrids comprise human/animal, animal/animal, or other such hybrid fVIII molecules, and further comprise at least one specific amino acid sequence in the A2 and/or C2 domain having one or more unique amino acids of the fVIII of one species substituted for the corresponding amino acid sequence (or amino acid) of the fVIII of the other species; or comprises at least one sequence in the A2 and/or C2 domain including one or more amino acids having no known sequence identity to fVIII substituted for specific amino acid sequence in human, animal, or hybrid fVIII. The resulting recombinant hybrid fVIII has reduced or no immunoreactivity to fVIII inhibitory antibodies, compared to human or porcine fVIII.

[0060] Unless otherwise specified or indicated, as used herein, "fVIII" denotes any functional fVIII protein molecule from human, any animal, any hybrid fVIII or any modified fVIII. "Hybrid fVIII" or "modified fVIII" denotes any functional fVIII protein, molecule or fragment thereof comprising fVIII amino acid sequences from one species substituted with one or more amino acids from another species, or with one or more amino acids having no known sequence identity with human or animal fVIII. Such hybrid and modified combinations include a fVIII amino acid sequence of human origin substituted with an amino acid sequence from an animal fVIII or an amino acid sequence having no known sequence identity to human or animal fVIII. Such combinations also include a fVIII amino sequence derived from more than two species, such as human/pig/mouse, or from one or more species in which an amino acid sequence having no known sequence identity to fVIII is substituted. Unless

otherwise indicated, "hybrid fVIII" and "modified fVIII" include fragments of the fVIII, which can be used as probes for research purposes or as diagnostic reagents.

[0061] A "fusion protein" or "fusion fVIII or fragment thereof", as used herein, is the product of a hybrid gene in which the coding sequence for one protein is extensively altered, for example, by fusing part of it to the coding sequence for a second protein from a different gene to produce a hybrid gene that encodes the fusion protein. As used herein, a fusion protein is a subset of the hybrid fVIII protein described in this application.

[0062] A "corresponding" nucleic acid or amino acid or sequence of either, as used herein, is one present at a site in a fVIII molecule or fragment thereof that has the same structure and/or function as a site in the fVIII molecule of another species, although the nucleic acid or amino acid number may not be identical. A DNA sequence "corresponding to" another fVIII sequence substantially corresponds to such sequence, and hybridizes to the sequence of the designated SEQ ID NO. under stringent conditions. A DNA sequence "corresponding to" another fVIII sequence also includes a sequence that results in the expression of a fVIII or fragment thereof and would hybridize to the designated SEQ ID NO. but for the redundancy of the genetic code.

[0063] A "unique" amino acid residue or sequence, as used herein, refers to an amino acid sequence or residue in the fVIII molecule of one species that is different from the homologous residue or sequence in the fVIII molecule of another species.

[0064] "Sequence homology" as used herein refers to identity or substantial similarity between two or more polypeptides or two or more nucleic acids. Sequence homology is determined on the basis of the nucleotide sequence of the two or more nucleic acids, or the amino acid sequence of the two or more polypeptides. The modified fVIII polypeptides of the present invention will have not more than 15%, usually not more than 5%, amino acid differences from the amino acid sequence recited in SEQ ID NO: 2, excluding the B domain.

[0065] "Specific activity," as used herein, refers to the activity that will correct the coagulation defect of human fVIII deficient plasma. Specific activity is measured in units of clotting activity per milligram total fVIII protein in a standard assay in which

the clotting time of human fVIII deficient plasma is compared to that of normal human plasma. One unit of fVIII activity is the activity present in one milliliter of normal human plasma. In the assay, the shorter the time for clot formation, the greater the activity of the fVIII being assayed. Porcine fVIII has coagulation activity in a human fVIII assay.

[0066] "Expression" refers to the set of processes that occur whereby genetic information is utilized to yield a product. For example, a DNA encoding the amino acid sequence of human fVIII having a number of amino acid substitutions can be "expressed" within a mammalian host cell to yield modified fVIII protein. The materials, genetic structures, host cells and conditions which permit expression of a given DNA sequence to occur are well-known in the art and can be manipulated to affect the time and amount of expression, as well as the intra- or extra-cellular location of the expressed protein. For example, by including DNA encoding a signal peptide at the 5' end of the DNA encoding porcine fVIII (the 5' end being, by convention, that end encoding the NH₂ terminus of the protein) the expressed protein becomes exported from the interior of the host cell into the culture medium. Providing a signal peptide coding DNA in combination with the modified fVIII coding DNA is advantageous because the expressed fVIII is exported into the culture medium which simplifies the process of purification. A preferred signal peptide is a mammalian fVIII signal peptide.

[0067] The human fVIII cDNA nucleotide sequence and predicted amino acid sequence are shown in SEQ ID NOs:1 and 2, respectively. FVIII is synthesized as an approximately 300 kDa single chain protein with internal sequence homology that defines the "domain" sequence NH₂-A1-A2-B-ap-A3-C1-C2-COOH. In a fVIII molecule, a "domain", as used herein, is a continuous sequence of amino acids that is defined by internal amino acid sequence identity and sites of proteolytic cleavage by thrombin. Unless otherwise specified, fVIII domains include the following amino acid residues, when the sequences are aligned with the human amino acid sequence (SEQ ID NO:2): A1, residues Ala1-Arg372; A2, residues Ser373-Arg740; B, residues Ser741-Arg1648; A3, residues Ser1690-Ile2032; C1, residues Arg2033-Asn2172; C2, residues Ser2173-Tyr2332. The remaining segment, residues Glu1649-Arg1689, is usually referred to as the fVIII light chain activation peptide. A

"B-domainless" or "B domain deleted" (BDD) fVIII, or fragment thereof, as used herein, refers to a fVIII protein that lacks part or the entire B domain.

[0068] FVIII is proteolytically activated by thrombin or factor Xa, which dissociates it from von Willebrand factor, forming fVIIIa, which has procoagulant function. The biological function of fVIIIa is to increase the catalytic efficiency of factor IXa toward factor X activation by several orders of magnitude. Thrombin-activated fVIIIa is a 160 kDa A1/A2/A3-C1-C2 heterotrimer that forms a complex with factor IXa and factor X on the surface of platelets or monocytes. A "partial domain" as used herein is a continuous sequence of amino acids forming part of a domain.

[0069] "Subunits" of human or animal fVIII, as used herein, are the heavy and light chains of the protein. The A3-C1-C2 domains, residues Ser1690-Tyr2332, comprise the fVIII light chain. The A1-A2 domains, residues 1-740, make up the fVIII heavy chain.

[0070] The terms "epitope," "antigenic site," and "antigenic determinant," as used herein, are used synonymously and are defined as a portion of the human, or animal fVIII, or fragments thereof, that is specifically recognized by an antibody. Lower antigenicity means antibodies, such as inhibitory antibodies, are less able to recognize or react to the site. An epitope (or antigenic site) can consist of any number of amino acid residues, and it can be dependent upon the primary, secondary, or tertiary structure of the protein.

[0071] The term "immunogenic site," as used herein, is defined as a region of the human or animal fVIII, or fragments thereof, that specifically elicits the production of an antibody to the fVIII, or fragment, in a human or animal, as measured by routine protocols, such as immunoassay, e.g. ELISA, or the Bethesda assay, described herein. It can consist of any number of amino acid residues, and it can be dependent upon the primary, secondary, or tertiary structure of the protein. Lower immunogenicity means antibodies, such as inhibitory antibodies, are less likely to be produced by the host animal or human. In some embodiments, the modified or hybrid fVIII equivalent or fragment thereof is non-immunogenic (does not elicit the production of antibodies) or has lower immunogenicity in an animal or human than human fVIII.

[0072] As used herein, a "hybrid fVIII equivalent molecule or fragment thereof" or "hybrid equivalent fragment fVIII or fragment thereof" is an active fVIII, a hybrid fVIII molecule or fragment thereof comprising at least one sequence having one or more amino acid residues, which have no known sequence identity to human or animal fVIII sequences, substituted for at least one sequence having one or more specific amino acid residues in the human, animal, or hybrid fVIII or fragment thereof. The sequence of one or more amino acid residues that have no known identity to human or animal fVIII sequence is also referred to herein as "non-fVIII amino acid sequence".

[0073] In an embodiment of the present invention, the amino acid(s), which are substituted into the fVIII and have no known sequence identity to human or animal fVIII, are alanine residues. In one embodiment, the specific fVIII sequence, for which the amino acid(s) having no known sequence identity to human or animal fVIII are substituted, includes an antigenic site that is immunoreactive with naturally occurring fVIII inhibitory antibodies, such that the resulting hybrid fVIII equivalent molecule or fragment thereof is less antigenic or not antigenic. In another embodiment, the specific fVIII sequence, for which the amino acid(s) having no known sequence identity to human or animal fVIII are substituted, includes an immunogenic site that elicits the formation of fVIII inhibitory antibodies in an animal or human, such that the resulting hybrid fVIII equivalent molecule or fragment thereof is less immunogenic or not immunogenic.

[0074] "fVIII deficiency," as used herein, includes deficiency in clotting activity caused by production of defective fVIII, by inadequate or no production of fVIII, or by partial or total inhibition of fVIII by inhibitors. Hemophilia A is a type of fVIII deficiency resulting from a defect in an X-linked gene and the absence or deficiency of the fVIII protein it encodes.

[0075] As used herein, "diagnostic assays" include assays that in some manner utilize the antigen-antibody interaction to detect and/or quantify the amount of a particular antibody that is present in a test sample to assist in the selection of medical therapies. There are many such assays known to those of skill in the art. As used herein, human, animal or modified human fVIII DNA, or fragment thereof, and protein expressed therefrom, in whole or in part, can be substituted for the

corresponding reagents in the otherwise known assays, whereby the modified assays may be used to detect and/or quantify antibodies to fVIII. It is the use of these reagents, the fVIII DNA, or fragment thereof, or protein expressed therefrom, that permits modification of known assays for detection of antibodies to human or animal fVIII. Such assays include, but are not limited to ELISAs, immunodiffusion assays, and immunoblots. Suitable methods for practicing any of these assays are known to those of skill in the art. As used herein, the fVIII or fragment thereof that includes at least one epitope of the protein can be used as the diagnostic reagent. Examples of other assays in which human, animal, such as porcine, or modified human fVIII or fragment thereof can be used include the Bethesda assay and anticoagulation assays.

[0076] The term "DNA encoding a protein, such as human fVIII" means a polydeoxynucleic acid whose nucleotide sequence embodies coding information to a host cell for the amino acid sequence of the protein, e.g. human fVIII, according to the known relationships of the genetic code.

[0077] The "expression product" of a DNA encoding a human or animal fVIII or a modified fVIII is the product obtained from expression of the referenced DNA in a suitable host cell, including such features of pre- or post-translational modification of protein encoded by the referenced DNA, including but not limited to glycosylation, proteolytic cleavage and the like. It is known in the art that such modifications can occur and can differ somewhat depending upon host cell type and other factors, and can result in molecular isoforms of the product, with retention of procoagulant activity. See, e.g. Lind, P. et al., *Eur. J. Biochem.* 232:1927 (1995), incorporated herein by reference.

[0078] An "expression vector" is a DNA element, often of circular structure, having the ability to replicate autonomously in a desired host cell, or to integrate into a host cell genome and also possessing certain well-known features which permit expression of a coding DNA inserted into the vector sequence at the proper site and in proper orientation. Such features can include, but are not limited to, one or more promoter sequences to direct transcription initiation of the coding DNA and other DNA elements such as enhancers, polyadenylation sites and the like, all as well known in the art. The term "expression vector" is used to denote both a vector

having a DNA coding sequence to be expressed inserted within its sequence, and a vector having the requisite expression control elements so arranged with respect to an insertion site that it can serve to express any coding DNA inserted into the site, all as well-known in the art. Thus, for example, a vector lacking a promoter can become an expression vector by the insertion of a promoter combined with a coding DNA.

GENERAL DESCRIPTION OF METHODS

[0079] The human fVIII gene was isolated and expressed in mammalian cells, as reported by Toole, J.J. et al. (1984) *Nature* 312:342-347; Gitschier, J. et al. (1984) *Nature* 312:326-330; Wood, W.I. et al. (1984) *Nature* 312:330-337; Vehar, G.A. et al. (1984) *Nature* 312:337-342; WO 87/04187; WO 88/08035; WO 88/03558; U.S. Patent No. 4,757,006, and the amino acid sequence was deduced from cDNA. U.S. Patent No. 4,965,199 to Capon et al. discloses a recombinant DNA method for producing fVIII in mammalian host cells and purification of human fVIII. Human fVIII expression on CHO (Chinese hamster ovary) cells and BHKC (baby hamster kidney cells) has been reported. Human fVIII has been modified to delete part or the entire B domain (U.S. Patent No. 4,868,112), and replacement of the human fVIII B domain with the human factor V B domain has also been attempted (U.S. Patent No. 5,004,803).

[0080] The cDNA sequence encoding human fVIII and predicted amino acid sequence are shown in SEQ ID NOs: 1 and 2, respectively. In SEQ ID NO: 1, the coding region begins at nucleotide position 208, the triplet GCC being the codon for amino acid number 1 (Ala) of the mature protein as given in SEQ ID NO: 2.

[0081] Human fVIII is isolated from plasma as a two subunit protein. The subunits, known as the heavy chain and light chain, are held together by a non-covalent bond that requires calcium or other divalent metal ions. The heavy chain of fVIII contains the A1 and A2 domains. The light chain of fVIII contains the A3, C1, and C2 domains. The B domain has no known biological function and can be removed, or partially removed, from the molecule proteolytically or by recombinant DNA technology methods without significant alteration in any measurable parameter

of fVIII. Human recombinant fVIII has a similar structure and function to plasma-derived fVIII, though it is not glycosylated unless expressed in mammalian cells.

[0082] Human activated fVIII ("fVIIIa") has three subunits due to cleavage of the heavy chain between the A1 and A2 domains. This structure is designated A1/A2/A3-C1-C2. Human fVIIIa is not stable under the conditions that stabilize porcine fVIIIa, presumably because of the weaker association of the A2 subunit of human fVIIIa. Dissociation of the A2 subunit of human and porcine fVIIIa is associated with loss of activity in the fVIIIa molecule. Yakhyæv, A. et al., 1997, *Blood* 90:Suppl. 1, Abstract #126, reported binding of A2 domain by low density lipoprotein receptor-related protein, suggesting that cellular uptake of A2 mediated by such binding acts to down-regulate fVIII activity.

[0083] Previous U.S. patents have disclosed how to modify human fVIII with porcine fVIII. U.S. patents 5,663,060 and 5,364,771 describe hybrid human/animal, particularly human/porcine, fVIII molecules having coagulant activity, in which elements of the fVIII molecule of human or an animal are substituted for corresponding elements of the fVIII molecule of the other species. The cDNA sequence encoding the complete A2 domain of porcine fVIII, predicted amino acid sequence, and hybrid human/porcine fVIII having substitutions of all domains, all subunits, and specific amino acid sequences were disclosed in U.S. Patent 5,364,771 and in WO 93/20093. More recently, the nucleotide and corresponding amino acid sequences of part of the A1 domain lacking the first 198 amino acid and of the A2 domain of porcine fVIII were reported in WO 94/11503, published May 26, 1994. The entire nucleotide sequence encoding porcine fVIII, including the complete A1 domain, activation peptide, A3, C1 and C2 domains, as well as the encoded amino acid sequence, was finally obtained by Lollar, as disclosed in U.S. Patent 5,859,204, issued January 12, 1999, and in WO 97/49725, published December 31, 1997, both incorporated herein by reference. The amino acid sequence of porcine fVIII is given in SEQ ID NO: 3.

[0084] Since current information indicates that the B domain has no inhibitory epitope and has no known effect on fVIII function, in some embodiments the B domain is wholly or partially deleted in the active hybrid or hybrid equivalent fVIII molecules or fragments thereof prepared by any of the methods described herein.

Expression of "B-domainless fVIII" is enhanced by including portions of the B-domain. The inclusion of the 3 amino acids of the B domain N-terminus and 11 amino acids of the B domain C-terminus was reported to result in favorable expression (Lind et al., 1995, *Eur. J. Biochem.* 232:19-27). "HBD" is a human fVIII that lacks the entire human B domain except for 5 amino acids of the B domain N-terminus and 9 amino acids of the B domain C-terminus (which results in the same 14 amino acid B domain linker sequence, S F S Q N P P V L K R H Q R, as disclosed in Lind et al.).

[0085] The purified modified fVIII or fragment thereof can be assayed for immunoreactivity and coagulation activity by standard assays including, for example, the plasma-free fVIII assay, the one-stage clotting assay, and the enzyme-linked immunosorbent assay using purified recombinant human fVIII as a standard.

[0086] Other vectors, including both plasmid and eukaryotic viral vectors, may be used to express a recombinant gene construct in eukaryotic cells depending on the preference and judgment of the skilled practitioner (see, for example, Sambrook et al., Chapter 16). Other vectors and expression systems, including bacterial, yeast, and insect cell systems, can be used but are not preferred due to differences in, or lack of, glycosylation.

[0087] Recombinant fVIII protein can be expressed in a variety of cells commonly used for culture and recombinant mammalian protein expression. In particular, a number of rodent cell lines have been found to be especially useful hosts for expression of large proteins. Preferred cell lines, available from the American Type Culture Collection, Rockville, MD, include baby hamster kidney cells, and Chinese hamster ovary (CHO) cells which are cultured using routine procedures and media.

FVIII molecules with reduced immunoreactivity:

[0088] Epitopes that are immunoreactive with antibodies that inhibit the coagulant activity of fVIII ("inhibitors" or "inhibitory antibodies") have been characterized based on known structure-function relationships in fVIII. Most inhibitory antibodies to human fVIII act by binding to epitopes located in the 40 kDa A2 domain or 20 kDa C2 domain of fVIII, disrupting specific functions associated with these domains, as described by Fulcher et al. (1985) *Proc. Natl. Acad. Sci USA* 82:7728-7732; and

Scandella et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:6152-6156. In addition to the A2 and C2 epitopes, there may be a third epitope in the A3 or C1 domain of the light chain of fVIII, according to Scandella et al. (1993) *Blood* 82:1767-1775. The significance of this putative third epitope is unknown, but it appears to account for a minor fraction of the epitope reactivity in fVIII.

[0089] After identification of clinically significant epitopes, recombinant fVIII molecules can be expressed that have less than or equal cross-reactivity compared with plasma-derived human or porcine fVIII when tested *in vitro* against a broad survey of inhibitor plasmas. Additional mutagenesis in epitopic regions can be done to reduce cross-reactivity. Reduced cross-reactivity, although desirable, is not necessary to produce a product that may have advantages over the existing plasma-derived human or porcine fVIII concentrate, which can produce side effects due to contaminant proteins or contaminant infectious agents such as viruses or prions. A recombinant fVIII or a modified recombinant fVIII molecule will not contain foreign proteins.

[0090] The basis for the greater coagulant activity of porcine fVIII appears to be the more rapid spontaneous dissociation of the human A2 subunit from human fVIIIa than the porcine A2 subunit from porcine fVIIIa. Dissociation of the A2 subunit leads to loss of activity, (Lollar, P. et al. (1990) *J. Biol. Chem.* 265:1688-1692; Lollar, P. et al. (1992) *J. Biol. Chem.* 267:23652-23657; Fay, P.J. et al. (1992) *J. Biol. Chem.* 267:13246-13250).

[0091] Anti-A2 antibodies block factor X activation, as shown by Lollar et al. (1994) *J. Clin. Invest.* 93:2497-2504. Previous mapping studies by deletion mutagenesis described by Ware et al. (1992) *Blood Coagul. Fibrinolysis* 3:703-716, located the A2 epitope to within a 20 kDa region of the NH₂-terminal end of the 40 kDa A2 domain. Competition immunoradiometric assays have indicated that A2 inhibitors recognize either a common epitope or narrowly clustered epitopes, as described by Scandella et al. (1992) *Throm. Haemostas.* 67:665-671, and as demonstrated in U.S. Patent 5,859,204.

[0092] Hemophilia A inhibitor patients and patients with acquired hemophilia A recognize immunodominant epitopes in the A2 and C2 domains of human fVIII.

Hemophilia A mice also recognize A2 and C2 domain epitopes when immunized with human fVIII using a dosing schedule that mimics use in human hemophilia A. The immune response of hemophilia A mice to human and porcine fVIII was compared using a domain specific ELISA. In this assay, monoclonal antibodies were tested against a panel of six single human fVIII domain hybrid human/porcine fVIII molecules as antigens that contain the human A1, A2, ap, A3, C1 or C2 domains. With anti-human antibodies, a positive signal with one of the single human domain proteins identifies domain specificity, whereas loss of signal indicates domain specificity of anti-porcine fVIII antibodies. Exon 16 (E16) – disrupted hemophilia A mice (n = 3) received six weekly 10 µg/kg intravenous injections of recombinant B-domain deleted human fVIII and a final 25 µg/kg boost. To obtain comparable inhibitor titers, E16 mice (n = 3) received six weekly injections of 40 µg/kg of recombinant B-domain deleted porcine fVIII. Spleens from high titer mice were fused with NS1 mouse myeloma cells and 485 of the resulting hybridomas were analyzed (Table 1).

Table 1: Domain Specificity

Mouse Immunogen Hybridomas			A1	A2	ap	A3	C1	C2	CR	MD
A	Human fVIII	95	2	16	0	2	7	21	23	24
B	Human fVIII	126	13	23	0	1	2	27	39	21
C	Human fVIII	54	1	15	1	2	1	10	9	15
D	Porcine fVIII	123	39	7	1	19	8	16	33	0
E	Porcine fVIII	27	13	5	0	0	0	4	2	3
F	Porcine fVIII	60	9	6	0	12	1	9	13	10

CR: Cross-reactive

MD: Multidomain

[0093] Human fVIII elicited a significantly greater number of antibodies to the A2 domain, whereas porcine fVIII elicited a significantly greater number of antibodies to the A1 and A3 domains ($p < 0.01$, chi square test). The greater number of anti-C2 antibodies to human fVIII was not statistically significant ($p = 0.10$) in this experiment. The differential immunodominance of human and porcine fVIII epitopes suggests that it may be possible to design a recombinant hybrid human/porcine fVIII molecule

that is less immunogenic than human fVIII in the treatment of patients with hemophilia A.

[0094] A triple mutant, R484A/R489A/P492A, which substitutes three alanine residues in the A2 epitope of a human B domain deleted fVIII (HBD), was found to be significantly less immunogenic than the unsubstituted B domain deleted human fVIII in hemophilia A mice (Fig. 4), while retaining full procoagulant activity. In contrast, the double mutant R484A/R489A, which substitutes two alanine residues in a B domain deleted human fVIII, was not less immunogenic than HBD. The clearance and hemostatic efficacy of R484A/R489A/P492A was similar to HBD (Fig. 5 and Table 2), making it unlikely the reduced immunogenicity of R484A/R489A/P492A was due to decreased bioavailability.

[0095] The decreased immunogenicity of R484A/R489A/P492A was identified by a reduction in the Bethesda inhibitor titer. In addition to inhibitory anti-A2 antibodies, the Bethesda assay can detect loss of fVIII coagulant function due to anti-C2 and other antibodies. In patient plasmas, anti-A2 and anti-C2 antibodies appear to contribute similarly to the inhibitor titer. In a previous study of 34 hemophilia A inhibitor patient plasmas, inhibitor titers were reduced by an average of 40% and 30% by soluble recombinant A2 and C2 domains, respectively (Prescott et al. (1997) *Blood* 89:3663-3671). The results using the R484A/R489A/P492A mutant were consistent with this previous study in that mutagenesis of A2 epitope resulted in a partial reduction of the inhibitor titer. Preliminary characterizations by homolog scanning mutagenesis of the B cell epitopes recognized by inhibitory antibodies in the murine model indicates that the A2 and C2 domains are the most frequently targeted domains as they are in humans. This result establishes the mouse model as a reliable predictive model of human efficacy. A combination of mutations in the A2 and C2 domains could further reduce the immunogenicity of human fVIII.

[0096] In contrast to the Bethesda assays, there was no significant difference among the treatment groups by anti-fVIII ELISA, which detects both inhibitory and non-inhibitory antibodies. Non-inhibitory antibodies and anti-fVIII antibodies have been identified in human hemophilia A patients (Giles et al., (1993) *Blood* 82:2452-2461). The relative contribution of inhibitory antibodies to the total ELISA signal is not known in human or murine hemophilia A. However, the results of the present

invention indicate that the ELISA assay is not sensitive to the reduction of immunogenicity of inhibitor epitopes because non-inhibitory antibodies also contribute to the total ELISA signal.

[0097] The immunodominance of the A2 epitope in hemophilia A mice was tested in plasmas from mice immunized with human B domainless fVIII using hybrid human/porcine and human/murine fVIII constructs. HP9, which is human except for insertion of porcine sequence within the 484 -508 segment of the A2 domain, was less antigenic than HBD (Fig. 2A). In contrast, HM1, which is murine except for insertion of human sequence within the 484 -508 segment, is more antigenic than murine B domainless fVIII (MBD) (Fig. 2B). Furthermore, there is a correlation between the reduction in antigenicity of HP9 compared to HBD and the increase in antigenicity of HM1 compared to MBD (Fig. 3) indicating that the antigenicity of the A2 epitope in individual mice is detected similarly in both assay systems.

[0098] A2 and C2 domain epitopes are immunodominant in fVIII inhibitor patients regardless of whether antibodies arise in the disparate immunological settings of alloimmunity or autoimmunity (Prescott et al., (1997) *Blood* 89:3663-3671). The R484 - I508 A2 segment appears to encompass the only inhibitory A2 epitope recognized by most patients (Healey et al., (1995) *J. Biol. Chem.* 270:14505-14509). This segment is predicted to consist of a large, surface exposed loop based on the homology model of the fVIII A domains (Pemberton et al., (1997) *Blood* 89:2413-2421). Inhibitory anti-C2 antibodies recognize a discontinuous epitope that contains a functionally important phospholipid binding site (Barrow et al., (2001) *Blood* 97:169-174; Arai et al., (1989) *J. Clin. Invest.* 83:1978-1984; Pratt et al., (1999) *Nature* 402:439-442; Spiegel et al., (2001) *Blood* 98:13-19). Although epitope-specific frequency distributions of the antibody populations in fVIII inhibitor patients or hemophilia A mice have not been enumerated, the B cell response to fVIII appears more restricted than the T cell response. For example, antibodies to the A1 domain are rarely seen, whereas the population of fVIII-specific T cells recognizes all of the domains in the human (Reding et al., (2000) *Thromb. Haemost.* 84:643-652) and murine (Wu et al., (2001)) *Thromb. Haemost.* 85:125-133) inhibitor response. Additionally, in contrast to the presence of common immunodominant B cell epitopes, immunodominant T cell epitopes have not been identified.

[0099] It has been established herein that structural modification of the fVIII molecule can reduce the immunogenicity in a murine hemophilia A model. The murine model can guide preclinical development of a therapeutically useful low immunogenic form of fVIII for human therapy. In one embodiment of the present invention, a method of identifying a modified fVIII having reduced immunogenicity or antigenicity is provided. This method comprises the steps: injecting at least one dose of said modified fVIII into a first group of animal (such as mice) test subjects; injecting at least one dose of an unmodified fVIII or a fVIII with known antigenic and/or immunogenic properties into a second group of animal test subjects; and using a diagnostic assay to compare the inhibitory antibodies produced by said first group of animal test subjects with inhibitory antibodies produced by said second group of animal test subjects.

[00100] Three human B domain deleted fVIII constructs, A2C2epi1, A2C2epi2 and A2C2epi3, contain the R484A/R489A/P492 mutation in the A2 domain and additional C2 domain mutations. In an attempt to conserve function, amino acids were selected for replacement of human residues at loops 1 and 3 (Fig. 6) in the C2 domain based on the sequences of other species for which the fVIII is known (Table 3). These three constructs were compared with human B domain deleted fVIII (HBD), R484A/R489A/ P492A (A2epi7), and each other.

[00101] A2C2epi1 contained the additional mutations M2199I/F2200L/L2252F in the C2 domain.

[00102] A2C2epi2 contained the additional mutations M2199I/F2200L/L2251V/L2252F in the C2 domain.

[00103] A2C2epi3 contained the additional mutations M2199L/F2200L/L2251V/L2252F in the C2 domain. A2C2epi3 differs from A2C2epi2 in that A2C2epi3 substitutes leucine for M2199 instead of isoleucine.

[00104] The five constructs, HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3, were expressed in BHK-derived cells and purified. The specific activities of A2C2epi1, A2C2epi2 and C3epi3 were similar to HBD (Table 4) and the purity of all the preparations was considered acceptable (Fig. 8).

[00105] A2C2epi3 was less immunogenic than HBD and A2epi7 (Figs. 9 and 10). In the Bethesda assay, A2C2epi1 was less immunogenic than HBD but not A2epi7. A2C2epi2 was not significantly less immunogenic than HBD in either assay, even though it contains the low immunogenicity R484A/R489A/P492A mutation. Conceivably, the C2 domain of A2C2epi2 is more immunogenic than the HBD, with the C2 domain offsetting a reduction of the immunogenicity of the A2 domain.

Diagnostic Assays

[00106] The fVIII cDNA and/or protein expressed therefrom, in whole or in part, can be used in assays as diagnostic reagents for the detection of inhibitory antibodies to human or animal fVIII or modified animal VIII in substrates, including, for example, samples of serum and body fluids of human patients with fVIII deficiency. These antibody assays include assays such as ELISA assays, immunoblots, radioimmunoassays, immunodiffusion assays, and assay of fVIII biological activity (e.g., by coagulation assay). Techniques for preparing these reagents and methods for use thereof are known to those skilled in the art.

[00107] Nucleic acid and amino acid probes can be prepared based on the sequence of the modified fVIII cDNA or protein molecule or fragments thereof. In some embodiments, these can be labeled using dyes or enzymatic, fluorescent, chemiluminescent, or radioactive labels that are commercially available. The amino acid probes can be used, for example, to screen sera or other body fluids where the presence of inhibitors to human, animal, or hybrid human/animal fVIII is suspected. Levels of inhibitors can be quantitated in patients and compared to healthy controls, and can be used, for example, to determine whether a patient with a fVIII deficiency can be treated with an animal or modified animal fVIII. The cDNA probes can be used, for example, for research purposes in screening DNA libraries.

Preparation of Recombinant FVIII

[00108] Recombinant fVIII can be produced through the use of eukaryotic protein expression systems. In general, a eukaryotic cell line, which is deficient in a required gene, is transformed with a vector comprising the gene that it has a deficiency for, and the recombinant DNA which one wishes to express. Transformation can be accomplished by techniques such as electroporation or viral delivery. The cell line

chosen to produce the protein is selected to be compatible with the protein of interest, capable of continuously expressing the protein of interests, capable of growing on a medium, which facilitates purification of the protein of interest, along with other factors known to those skilled in the art. Examples of such techniques are disclosed in European Patent Application 0 302 968 A2 and United States Patent No. 5,149,637 both of which are incorporated by reference in their entirety.

Testing of Recombinant FVIII Molecules

[00109] The recombinant fVIII molecules can be tested in humans for their reduced antigenicity and/or immunogenicity in at least two types of clinical trials. In one type of trial, designed to determine whether the recombinant or recombinant hybrid fVIII is antigenic with inhibitory antibodies, recombinant fVIII or recombinant modified fVIII is administered, preferably by intravenous infusion, to approximately 25 patients having fVIII deficiency who have antibodies to fVIII that inhibit the coagulant activity of therapeutic human or porcine fVIII. The dosage of the recombinant or recombinant modified fVIII is in a range between 5 and 50 Units/kg body weight, preferably 10-50 Units/kg, and most preferably 40 Units/kg body weight. Approximately 1 hour after each administration, the recovery of fVIII from blood samples is measured in a one-stage coagulation assay. Samples are taken again approximately 5 hours after infusion, and recovery is measured. Total recovery and the rate of disappearance of fVIII from the samples are predictive of the antibody titer and inhibitory activity. If the antibody titer is high, fVIII recovery usually cannot be measured. The recovery results are compared to the recovery results in patients treated with plasma-derived human fVIII, recombinant human fVIII, porcine fVIII, and other commonly used therapeutic forms of fVIII or fVIII substitutes.

[00110] In a second type of clinical trial, designed to determine whether the recombinant or recombinant modified fVIII is immunogenic, i.e., whether patients will develop inhibitory antibodies, recombinant or recombinant hybrid fVIII is administered, as described in the preceding paragraph, to approximately 100 previously untreated hemophiliac patients who have not developed antibodies to fVIII. Treatments are given approximately every 2 weeks over a period of 6 months to 1 year. At 1 to 3 month intervals during this period, blood samples are drawn and Bethesda assays or other antibody assays are performed to determine the presence

of inhibitory antibodies. Recovery assays can also be done, as described above, after each infusion. Results are compared to hemophiliac patients who receive plasma-derived human fVIII, recombinant human fVIII, porcine fVIII, or other commonly used therapeutic forms of fVIII or fVIII substitutes.

Pharmaceutical Compositions

[00111] Pharmaceutical compositions comprising recombinant or recombinant modified fVIII, alone or in combination with appropriate pharmaceutical stabilization compounds, delivery vehicles, and/or carrier vehicles, are prepared according to known methods, as described in Remington's *Pharmaceutical Sciences* by E.W. Martin.

[00112] In one preferred embodiment, the preferred carriers or delivery vehicles for intravenous infusion are physiological saline or phosphate buffered saline.

[00113] In another preferred embodiment, suitable stabilization compounds, delivery vehicles, and carrier vehicles include but are not limited to other human or animal proteins such as albumin.

[00114] Phospholipid vesicles or liposomal suspensions are also preferred as pharmaceutically acceptable carriers or delivery vehicles. These can be prepared according to methods known to those skilled in the art and can contain, for example, phosphatidylserine/phosphatidylcholine or other compositions of phospholipids or detergents that together impart a negative charge to the surface, since fVIII binds to negatively charged phospholipid membranes. Liposomes may be prepared by dissolving appropriate lipid(s) (such as stearyl phosphatidyl ethanolamine, stearyl phosphatidyl choline, arachadoyl phosphatidyl choline, and cholesterol) in an inorganic solvent that is then evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the hybrid or modified fVIII is then introduced into the container. The container is then swirled by hand to free lipid material from the sides of the container and to disperse lipid aggregates, thereby forming the liposomal suspension.

[00115] Recombinant or recombinant modified fVIII can be combined with other suitable stabilization compounds, delivery vehicles, and/or carrier vehicles, including

vitamin K dependent clotting factors, tissue factor, and von Willebrand factor (vWf) or a fragment of vWf that contains the fVIII binding site, and polysaccharides such as sucrose.

[00116] Recombinant or recombinant modified fVIII can also be delivered by gene therapy in the same way that human fVIII can be delivered, using delivery means such as retroviral vectors. This method consists of incorporation of fVIII cDNA into human cells that are transplanted directly into a fVIII deficient patient or that are placed in an implantable device, permeable to the fVIII molecules but impermeable to cells, which are then transplanted. The preferred method will be retroviral-mediated gene transfer. In this method, an exogenous gene (e.g., a fVIII cDNA) is cloned into the genome of a modified retrovirus. The gene is inserted into the genome of the host cell by viral machinery where it will be expressed by the cell. The retroviral vector is modified so that it will not produce virus, preventing viral infection of the host. The general principles for this type of therapy are known to those skilled in the art and have been reviewed in the literature (e.g., Kohn, D.B. et al. (1989) *Transfusion* 29:812-820).

[00117] Recombinant or recombinant modified fVIII can be stored bound to vWf to increase the half-life and shelf-life of the modified molecule. Additionally, lyophilization of fVIII can improve the yields of active molecules in the presence of vWf. Current methods for storage of human and animal fVIII used by commercial suppliers can be employed for storage of hybrid fVIII. These methods include: (1) lyophilization of fVIII in a partially-purified state (as a fVIII "concentrate" that is infused without further purification); (2) immunoaffinity-purification of fVIII by the Zimmerman method and lyophilization in the presence of albumin, which stabilizes the fVIII; (3) lyophilization of recombinant fVIII in the presence of albumin.

[00118] Additionally, hybrid or modified fVIII has been indefinitely stable at 4° C in 0.6 M NaCl, 20 mM MES, and 5 mM CaCl₂ at pH 6.0 and also can be stored frozen in these buffers and thawed with minimal loss of activity.

Methods of Treatment

[00119] Recombinant or recombinant modified fVIII is used to treat uncontrolled bleeding due to fVIII deficiency (e.g., intraarticular, intracranial, or gastrointestinal

hemorrhage) in hemophiliacs with and without inhibitory antibodies and in patients with acquired fVIII deficiency due to the development of inhibitory antibodies. The active materials are preferably administered intravenously. It is especially useful to treat naïve patients, i.e., patients who have not yet developed inhibitory antibodies, with FVIII molecules having low immunogenicity in order to reduce the likelihood of the patient developing inhibitory antibodies.

[00120] Additionally, recombinant or recombinant modified fVIII can be administered by transplant of cells genetically engineered to produce the hybrid or by implantation of a device containing such cells, as described above.

[00121] In a preferred embodiment, pharmaceutical compositions of recombinant or recombinant modified fVIII alone or in combination with stabilizers, delivery vehicles, and/or carriers are infused into patients intravenously according to the same procedure that is used for infusion of human or animal fVIII.

[00122] The treatment dosages of recombinant or recombinant modified fVIII composition that must be administered to a patient in need of such treatment will vary depending on the severity of the fVIII deficiency. Generally, dosage level is adjusted in frequency, duration, and units in keeping with the severity and duration of each patient's bleeding episode. Accordingly, the modified fVIII is included in the pharmaceutically acceptable carrier, delivery vehicle, or stabilizer in an amount sufficient to deliver to a patient a therapeutically effective amount of the fVIII to stop bleeding, as measured by standard clotting assays.

[00123] FVIII is classically defined as that substance present in normal blood plasma that corrects the clotting defect in plasma derived from individuals with hemophilia A. The coagulant activity *in vitro* of purified and partially-purified forms of fVIII is used to calculate the dose of fVIII for infusions in human patients and is a reliable indicator of activity recovered from patient plasma and of correction of the *in vivo* bleeding defect. There are no reported discrepancies between standard assay of novel fVIII molecules *in vitro* and their behavior in the dog infusion model or in human patients, according to: Lusher, J.M. et al. 328 *New Engl. J. Med.* 328:453-459; Pittman, D.D. et al., (1992) *Blood* 79:389-397; and Brinkhous et al. (1985) *Proc. Natl. Acad. Sci.* 82:8752-8755.

[00124] Usually, the desired plasma fVIII level to be achieved in the patient through administration of the recombinant or recombinant modified fVIII is in the range of 30-100% of normal. In a preferred mode of administration of the recombinant or recombinant modified fVIII, the composition is given intravenously at a preferred dosage in the range from about 5 to 50 units/kg body weight, more preferably in a range of 10-50 units/kg body weight, and most preferably at a dosage of 20-40 units/kg body weight; the interval frequency is in the range from about 8 to 24 hours (in severely affected hemophiliacs); and the duration of treatment in days is in the range from 1 to 10 days or until the bleeding episode is resolved. See, e.g., Roberts, H.R., and M.R. Jones, Ch. 153, 1453-1474, 1460, in Hematology, Williams, W. J., et al., ed. (1990). Patients with inhibitors may require more recombinant or recombinant modified fVIII, or patients may require less recombinant or recombinant modified fVIII because of its higher specific activity than human fVIII or decreased antibody reactivity or immunogenicity. As in treatment with human or porcine fVIII, the amount of recombinant or recombinant modified fVIII infused is defined by the one-stage fVIII coagulation assay and, in selected instances, *in vivo* recovery is determined by measuring the fVIII in the patient's plasma after infusion. It is to be understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

[00125] Treatment can take the form of a single intravenous administration of the composition or periodic or continuous administration over an extended period of time, as required. Alternatively, recombinant or recombinant modified fVIII can be administered subcutaneously or orally with liposomes in one or several doses at varying intervals of time.

[00126] FVIII can also be used to treat uncontrolled bleeding due to fVIII deficiency in hemophiliacs who have developed antibodies to human fVIII. In this case, coagulant activity that is superior to that of human or animal fVIII alone is not necessary. Coagulant activity that is inferior to that of human fVIII (i.e., less than

3,000 units/mg) will be useful if that activity is not neutralized by antibodies in the patient's plasma.

[00127] The recombinant or recombinant modified fVIII molecule and the methods for isolation, characterization, making, and using it generally described above will be further understood with reference to the following non-limiting examples.

EXAMPLES

[00128] Example 1: Construction and evaluation of the R484A/R489A/P492A (A2epi7) mutant

[00129] *Materials* – Citrated hemophilia A plasma and normal pooled human plasma (FACT) were purchased from George King Biomedical, Inc. (Overland Park, KA). Activated partial thromboplastin time reagent (Automated APTT ®) was purchased from Biomerieux (Durham, NC). Murine anti-human fVIII monoclonal antibodies ESH4, ESH5 and ESH8 were purchased from American Diagnostica. Synthetic oligonucleotides were purchased from Life Technologies. Restriction enzymes were purchased from New England Biolabs or Promega. A cell line derived from baby hamster kidney cells was a generous gift from Dr. R.T.A. Macgillivray (Funk et al., 1990, *Biochemistry* 29:1654-1660.). Exon 16-disrupted (E16) hemophilia A mice in a C57BL/6 background were obtained from Dr. Leon Hoyer and a breeding colony was established (Bi et al., 1995, *Nat. Genet.* 10:119-121). Nine- to twelve-week old E16 male or female hem A or normal C57BL/6 mice were used in the experiments. Novel fVIII DNA sequences generated by PCR were confirmed by dideoxy sequencing using an Applied Biosystem 373a automated DNA sequencer and the PRISM dye terminator kit.

[00130] *Construction of recombinant fVIII mutant cDNAs* – The cDNA encoding a human B-domain deleted (HBD) form of fVIII was prepared as described in Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349. It contains a S F S Q N P P V L K R H Q R linker sequence between the A2 and *ap*-A3 domains. The linker corresponds to the first five and last nine amino acids of the B domain and contains a recognition sequence for intracellular PACE/furin processing. This produces A1-A2/*ap*-A3-C1-C2 heterodimeric fVIII as the dominant secreted species, which is considered the physiologic form. The cDNAs for porcine and murine B-domain deleted forms of

fVIII, designated PBD and MBD, respectively, which also contain PACE/furin recognition linker sequences, were prepared as described previously (Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349; Doering et al., 2002, *Thromb. Haemost.* 88:450-458). The cDNA for a B domainless hybrid/porcine fVIII molecule designated HP9, which contains insertion of the porcine segment corresponding to residues 484-508 in the A2 domain of human fVIII (Fig. 1), has been described previously (Healey et al., 1995, *J. Biol. Chem.* 270:14505-14509). A B domainless hybrid human/murine fVIII cDNA, HM1, encoding a fVIII molecule that is murine except for a human segment corresponding to residues 484-508 of the A2 domain (Fig. 1), was constructed by splicing-by-overlap extension mutagenesis using MBD as the template.

[00131] The construction of a cDNA encoding a B domainless R489A human fVIII has been described previously (Lubin et al., 1997, *J. Biol. Chem.* 272:30191-30195) and was modified further by insertion of DNA encoding the S F S Q N P P V L K R H Q R linker sequence. The resulting cDNA was used as a template for the production cDNAs encoding R484A/R489A and R484A/R489A/P492A human fVIII (Fig. 1) by splicing-by-overlap extension mutagenesis. For the R484A/R489A mutant, 5'- CAC GGA ATC ACT GAT GTC GCC CCT TTG TAT TCA GCC AGA -3' and 5'- TCT GGC TGA ATA CAA AGG GGC GAC ATC AGT GAT TCC GTG -3' were used as the mutagenic sense and antisense primers, respectively. For the R484A/R489A/P492A mutant 5'- ACT GAT GTC GCC CCT TTG TAT TCA GCC AGA TTA GCC AAA -3' and 5'- TTT GGC TAA TCT GGC TGA ATA CAA AGG GGC GAC ATC AGT -3', were used as the mutagenic sense and antisense primers, respectively.

[00132] *Expression and purification of recombinant fVIII molecules* – Recombinant fVIII molecules were expressed in baby hamster kidney – derived cells in serum-free medium using the ReNeo expression vector as described previously in Healey et al., 1998, *Blood* 92:3701-3709. HBD, R484A/R489A, R484A/R489A/P492A, MBD, PBD and HM1 were purified by SP-Sepharose Fast Flow and Source Q or Mono Q ion-exchange chromatography essentially as describe previously for HBD, MBD and PBD in Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349; and Doering et al., 2002, *Thromb. Haemost.* 88:450-458.

[00133] HP9 was purified using the following procedure. Ammonium sulfate was added to 5.9 liters of cell culture medium at 4°C to 65% saturation and allowed to stir overnight. The precipitate was collected by centrifugation, dialyzed against 0.15 M NaCl, 0.02 M Hepes, 5mM CaCl₂, 0.01% Tween-80, pH 7.4, and applied to a 1.5 X 10 cm ESH5 – Sepharose column equilibrated in the same buffer. HP9 was eluted with 1 M NaCl, 5 mM Mes, 2.5 mM CaCl₂, 50% ethylene glycol (v/v), pH 6.0. Fractions from the fVIII activity peak were diluted 1/5 into 0.04 M Hepes, 5 mM CaCl₂, 0.01% Tween-80, pH 7.4 and further purified using Mono Q ion-exchange chromatography.

[00134] Purified proteins were at least 90% pure as judged by sodium dodecyl sulfate – polyacrylamide gel electrophoresis and contained heterodimeric fVIII as the dominant species. Concentrations were calculated using the absorbance at 280 nm of the purified proteins and molar extinction coefficients that were estimated from the respective deduced amino acid compositions (Pace et al., 1995, *Protein Sci.* 4:2411-2423). The specific activities of the purified proteins were calculated using the protein concentration and fVIII coagulant activity, which was measured as described below against a human fVIII plasma standard. The following specific activities were obtained: HBD, 1300 U/nmole; R484A/R489A/P492A, 1270 U/mole; R484A/R489A/P492A, 1460 U/nmol; HP9 1100, U/nmol; HM1 540, U/nmol. The specific activities of the PBD and MBD preparations have been previously reported as 2050 U/nmol and 660 U/nmol, respectively (Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349; and Doering et al., 2002, *Thromb. Haemost.* 88:450-458). The deduced polypeptide chain molecular mass for all fVIII species used in this study is 165 kDa and was for conversion of fVIII concentrations to mass units.

[00135] *Immunization of hemophilia A mice with HBD, R484A/R489A/P492A and R484A/R489A*- Preparations of HBD, R484A/R489A/P492A and R484A/R489A were diluted to 0.17 mg/ml in 0.4 M NaCl, 20 mM HEPES, 5 mM CaCl₂ 0.01% Tween-80 pH 7.4 and stored in small aliquots at -80°C. Samples were diluted were diluted to 4 µg/ml in sterile normal saline immediately prior to injection. Eighty-three hemophilia A mice were divided into HBD (n=24), R484A/R489A/P492A (n=23), R484A/R489A (n=24), buffer injected control (n=5) and non-injected control (n=7) groups. Mice were warmed under a 75-watt lamp for 3 to 4 minutes to dilate tail veins before

injection. Silver nitrate was used to cauterize bleeding after injections and tail snips. Recipients of different test materials were mixed within each cage and remained in their original cages to avoid fighting. Approximately equal numbers of males and females were in each group. Mice received six injections of 10 µg/kg body weight (~80 U/kg) at 14 day intervals, followed by a final rejection of 25 µg/kg body weight (~200 U/kg) two weeks after the sixth dose. Blood was collected into 1/10 volume 3.8% trisodium citrate by tail snip under metofane anesthesia 13 days after the fourth injection and by terminal cardiac puncture 13-14 days after the final injection. Samples were held on ice prior to centrifugation at 3000 x g for 15 minutes at 4°C to collect plasma. During the course of the experiment 0, 4, 5, 1 and 2 mice died in the HBD, R484A/R489A/P492A, R484A/R489A, buffer injected control and non-injected control groups, respectively. The overall mortality during this experiment was 14%, which was considered acceptable given the increased mortality of hemophilia A mice associated with handling (Bi et al., 1995, *Nat. Genet.* 10:119-121).

[00136] *In vivo clearance of HBD and R484A/R489A/P492A* – For clearance studies, HBD or R484A/R489A/P492A were diluted to 50 U/ml (~6.25 µg/ml) in sterile saline for injection immediately before use. Hemophilia A mice were anesthetized with metofane, weighed, warmed under a 75-watt lamp for 3 minutes to dilate veins and injected with 100 U/kg (~12.5 µg/kg) by tail vein. Silver nitrate was used to cauterize the injection sites. At various time, mice were anesthetized by intraperitoneal injection of a mixture of 300 mg/kg ketamine and 75 mg/kg xylazine and blood was collected by cardiac puncture into 1/10 volume of 3.8% trisodium citrate. Because serial sampling from hemophilia A mice is difficult, three mice were used for each time point for each fVIII construct. Plasma samples were prepared by centrifugation and stored at -70°C. FVIII activity was determined by chromogenic assay as described below.

[00137] *Hemostatic efficacy of HBD and R484A/R489A/P492A* – The hemostatic efficacy of R484A/R489A/P492A in hemophilia A mice was measured using a tail vein transection model as described previously (Parker et al., 2003, *Thromb. Haemost.* 89:480-485). Briefly, mice were anesthetized with 1.5 mg/kg droperidol/75 mg/kg ketamine intraperitoneally, warmed to dilate the tail veins, and injected with R484A/R489A/P492A. After anesthesia was deepened using methoxyflurane, mice

were placed in a 50 ml conical restraint tube, the distal 1 cm of tail was transected and the stump was placed in a test tube containing 150 mM NaCl at 37°C. At 2 h, surviving mice were caged and mortality at 24 h was determined. The up-and-down method for small samples (Dixon, W.J., 1965, *J. Amer. Stat. Assoc.* 60:967-978; and Dixon and Massey, 1969, in *Introduction to Statistical Analysis*, McGraw Hill, New York, 377-394) was used to estimate the dose that produces 50% survival (ED₅₀). An initial dose of fVIII was given to a single mouse as a *priori* estimate of the ED₅₀. If the mouse survived the 24 h test period, another mouse was tested and the dose was decreased. If the subject died, the dose was increased in the next subject. Testing was continued until a chosen nominal sample size of six was reached. A constant log dose increment or decrement of 0.1, corresponding to a dilution factor of 1.26 was used. The ED₅₀ was calculated using the equation:

$$\text{Log ED}_{50} = x_f + kd$$

Where x_f is the logarithm of final test dose, d is the log dose increment or decrement, and k is obtained from a table based on maximum likelihood estimates (Dixon, W.J., 1965, *J. Amer. Stat. Assoc.* 60:967-978). The standard error of log ED₅₀ (s.e.) was estimated using the equation :

$$\text{s.e.} = \sigma \sqrt{a}$$

where a equals 0.31 for a nominal sample size of six (Dixon, W.J., 1991 *Neuroscience & Biobehavioral Reviews* 15:47-50) and σ is the population standard deviation, which was assumed to be 0.12 (Bruce, R.D., 1985, *Fundamentals & Applied Toxicology* 5:151-157).

[00138] *FVIII coagulation assays and ELISAs* – FVIII activity was measured by one-stage clotting assay (Bowie and Owen, 1984, in Disorders of Hemostasis, O.D. Ratnoff and Forbes, C.D., editors, Grune & Stratton, Inc., Orlando, 43-72) as described previously (Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349) using normal human plasma (FACT) as the standard. FVIII inhibitor titers were measured by a modified Bethesda assay in which fVIII constructs (HBD, R484A/R489A, R484A/R489A/P492A, HP9 or PBD) were added to hemophilia A plasma to a final concentration of 0.8 – 1.2 units per ml incubated with varying concentrations of inhibitor for 2 hours at 37°C. One Bethesda unit (BU) is defined as the amount of inhibitory activity that produces 50% inhibition of fVIII activity in the one-stage clotting

assay. The 50% inhibition point was identified by interpolation using only data points falling within a range of 40-60% inhibition. An average of at least three data points in this range were used for each determination. Because of the smaller volumes required, fVIII activity was measured in the pharmacokinetic study by a chromogenic assay (Coamatic, Chromogenix/Diapharma Group, West Chester, OH) according to instructions supplied by the manufacturer.

[00139] Antibodies to HBD, R484A/R489A, R484A/R489A/P492A, MBD or HM1 were measured by sandwich ELISA by immobilizing the respective fVIII antigens and using ESH4 and biotinylated-ESH8 as capture and detection antibodies, respectively, as described previously (Lubin et al., 1994, *J. Biol. Chem.* 269:8639-8641). Absorbance values obtained from seven dilution of test plasma were plotted versus the logarithm of the plasma dilution and the resulting sigmoidal curves were fit to modified version of the 4-parameter logistic fit equation by nonlinear regression using the Levenberg-Marquardt algorithm. The ELISA titer was defined empirically as the dilution of plasma that returns an absorbance value of 0.3 derived from the fitted curves.

[00140] *Immunodominance of human R484-I508 epitope in hemophilia A mice* – To reduce the immunogenicity of human fVIII, two HBD fVIII mutants were developed that contain alanine substitutions of antigenic amino acids in the A2 domain. These mutants, designated R484A/R489A and R484A/R489A/P492A, were compared (Fig. 1) to “wild-type” HBD fVIII in hemophilia A mice using an intravenous immunization protocol described above. Two approaches were used to determine whether hemophilia A mice recognize the R484-I508 A2 sequence that is immunodominant in humans. First, plasmas from the HBD treatment group were tested for reactivity against a hybrid human/porcine fVIII molecule, designated HP9, which is human except for insertion of porcine sequence within the 484-508 segment (Fig. 1). If immune plasmas contain antibodies that recognize the R484-I508 epitope, they should react less well with the HP9 molecule than HBD because of the incomplete cross-reactivity of fVIII inhibitors with human and porcine fVIII. The average Bethesda titers in the HBD treatment group against HBD and B domainless porcine fVIII were 680 and 31 Bethesda units, respectively, corresponding to a cross-reactivity of 4%, demonstrating that porcine fVIII is poorly cross-reactive in this

model. Figure 2A shows paired data comparing the inhibition of HBD and HP9 by individual mouse plasmas in the Bethesda assay. Bethesda titers of most of the pairs were decreased using HP9 as the target antigen compared to HBD. The difference between HP9 and HBD groups was statistically significant ($p < 0.0001$, paired t test). Figure 2A also shows that all of the plasmas recognized HP9, indicating the presence of inhibitory epitopes directed toward human outside of the R484-I508 A2 epitope.

[00141] The immunodominance of the human R484-I508 segment in hemophilia A mice also was tested by a converse experiment in which the same plasma were tested by ELISA for reactivity against a hybrid human/murine fVIII molecule, designated HM1, which is murine except for human R484-I508 segment (Fig. 1). The average ELISA titer against B domainless murine fVIII, MBD, was 19% of that against HBD, demonstrating that plasmas cross-react poorly with murine fVIII, and thus that human/murine hybrid fVIII molecules can be used to study differential antigenicity. The HM1 hybrid would be expected to be more antigenic compared to MBD because of the presence of the antigenic human R484-I508 segment. Figure 2B shows that ELISA titers of most of the plasmas were increased using HM1 as the target antigen compared to MBD. The difference between the HM1 and MBD groups was statistically significant ($p < 0.02$, paired t test).

[00142] Additionally, in the plasmas from mice immunized with HBD, there was a significant correlation between the reduction in antigenicity of HP9 compared to HBD and the increase in antigenicity of HM1 compared to MBD (Fig. 3). Thus, the degree to which individual mice developed an immune response to the R484-I508 epitope could be detected similarly in both assay systems. This indicates that the reduced antigenicity of HP9 and the increased antigenicity are not artifactual, for example, due to differences in antigen binding to microtiter plates.

[00143] *Comparative immunogenicity of HBD, R484A/R489A/P492A, R484A/R489A* – The hemophilia A mice in the HBD, R484A/R489A, and R484A/R489A/P492A treatment groups received six intravenous injections of 10 $\mu\text{g/kg}$ at 14 day intervals and a final injection of 25 $\mu\text{g/kg}$ body as described in example 1. Plasmas obtained two weeks after the last injection were tested for anti-fVIII antibodies by Bethesda assay and by ELISA. Figure 4 shows that inhibitor titers

were lower in the R484A/ R489A/P492A group ($p= 0.01$, Mann-Whitney U test). In contrast, inhibitor levels in the R484A/ R489A were not significantly different from the HBD group. Plasmas in the three groups also were tested for anti-fVIII antibodies by ELISA. There was no significant difference between the groups.

[00144] *In vivo clearance of HBD and R484A/R489A/P492A* – The immune response following intravenous injection depends on delivery of the immunogen to the spleen. Thus, instead of being intrinsically less immunogenic than HBD, R484A/ R489A/P492A can be cleared more rapidly from the circulation. Figure 5 shows that the pharmacokinetics of R484A/ R489A/P492A and HBD are similar in hemophilia A mice.

[00145] *Comparative efficacy of HBD and R484A/ R489A/P492A* – The hemostatic efficacy of R484A/ R489A/P492A was measured in hemophilia A mice using a tail vein transection model (Parker and Lollar, (2003) *Thromb. Haemost.* 89:480-485). The estimated dose of R484A/ R489A/P492A that produces 50% survival (ED_{50}) was measured using the up-and-down method (Dixon, W.J., (1965) *J. Amer. Stat. Assoc.* 60:967-978; Dixon and Massey, (1969) "Sensitivity Experiments," in Introduction to Statistical Analysis, McGraw Hill, New York, 377-394). The results are shown in Table 2. The method yielded an ED_{50} of 47.5 units/kg (95% confidence interval, 34.9 – 64.6 units/kg) for R484A/R489A/P492A. For HBD, and ED_{50} of 57.7 units/kg (95% confidence interval, 42.4 – 78.5 units/kg) has been published (Parker and Lollar, (2003) *Thromb. Haemost.* 89:480-485). The published data for HBD and R484A/ R489A/P492A data in the current study were collected contemporaneously on matched littermates. The comparison indicates that the hemostatic efficacy of HBD and R484A/ R489A/P492A in this model are indistinguishable.

Table 2: Hemostatic efficacy of R484A/R489A/P492A in hemophilia A mice

Dose (units/kg)	Day 0	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6
39.8			X				X
50.1		O		X		O	
63.1	O				O		

$ED_{50} = 47.5$ units/kg*

95% confidence interval: 34.9 – 64.6 units/kg*

O - Alive X - Dead

[00146] The E16 knockout hemophilia A mice used in this study, along with the genetically similar, phenotypically indistinguishable E17 knockout mice, contained targeted disruptions in the region of the fVIII gene that encodes the A3 domain (Bi et al., (1995) *Nat. Genet.* 10:119-121; Bi et al., (1996) *Blood* 88:3446-3450). Both strains secrete a non-functional polypeptide containing sequence NH2-terminal to A3 domain, including the A2 domain (Sarkar et al., (2000) *Hum. Gene Ther.* 11:881-894). In contrast to the murine hemophilia A mice that are available, human hemophilia A is extremely heterogeneous (Tuddenham et al., (1994) *Nucleic. Acids. Res.* 22:4851-4868). Mutations produce phenotypes that range from no detectable synthesis of fVIII to secretion of partly to completely nonfunctional fVIII. E16 and E17 mice are analogous to hemophilia A patients who secrete nonfunctional fVIII. The E16 mice are not tolerant to the A2 domain despite the presence of truncated circulating fVIII that contains the A2 domain (Figs. 2 and 3). This may be due to non-native folding of the truncated fVIII polypeptide chain, which prevents induction of tolerance.

[00147] An immunogenicity model was used in which fVIII is given intravenously to hemophilia A mice using a dosage schedule that mimics the use of fVIII in human hemophilia A (Qian et al., (1999) *Thromb. Haemost.* 81:240-244). In this model, hemophilia A mice develop a T cell-dependent antibody response (Qian et al., (1999) *Thromb. Haemost.* 81:240-244; Wu et al., (2001) *Thromb. Haemost.* 85:125-133; Reipert et al., (2000) *Thromb. Haemost.* 84:826-832) that can be inhibited by blockade of the CD28-B7 (Qian et al., (2000) *Blood* 95:1324-1329) or CD40-CD154 (Qian et al., *Eur. J. Immunol.* 30:2548-2554; Rossi et al., (2001) *Blood* 97:2750-2757; Reipert et al., (2002) *Thromb. Haemost.* 86:1345-1352) co-stimulation pathways. The anti-fVIII antibody response is not isotypically restricted and its distribution is not different from normal mice. (Wu et al., (2001) *Thromb. Haemost.* 85:125-133; Reipert et al., (2000) *Thromb. Haemost.* 84:826-832). The effects on immunogenicity observed with the recombinant or recombinant modified fVIII molecules described herein are therefore likely to be comparative to those observed in humans.

Example 2: Construction and evaluation of the A2C2epi1, A2C2epi2 and A2C2epi3 mutants

[00148] *Construction of recombinant fVIII mutant cDNAs* – The cDNA encoding a human B-domain deleted (HBD) form of fVIII was prepared as described in Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349. It contains a S F S Q N P P V L K R H Q R linker sequence between the A2 and ap-A3 domains. The A2epi7 cDNA was prepared as described in Example 1. The A2C2epi1, A2C2epi2 and A2C2epi3 cDNAs were prepared by splicing-by-overlap extension mutagenesis using A2epi7 as a template.

[00149] *Expression and purification of recombinant fVIII molecules* – Recombinant fVIII molecules were expressed in baby hamster kidney – derived cells in serum-free medium using the ReNeo expression vector as described previously in Healey et al., 1998, *Blood* 92:3701-3709. HBD and A2epi7 were isolated using SP-Sepharose Fast Flow and Source Q ion-exchange chromatography essentially as describe previously for HBD in Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349; and Doering et al., 2002, *Thromb. Haemost.* 88:450-458. Because of the relatively low yields of A2C2epi1, A2C2epi2 and A2C2epi3 in cell culture, an additional monoS chromatography step was used.

[00150] *Immunization of hemophilia A mice* – FVIII preparations were diluted to 0.17 mg/ml in 0.4 M NaCl, 20 mM HEPES, 5 mM CaCl₂, 0.01% Tween-80 pH 7.4 and stored in small aliquots at -80°C. Samples were diluted were diluted to 4 µg/ml in sterile normal saline immediately prior to injection. Nine- to twelve-week old E16 male or female hemophilia A mice were randomized to five groups of 125. Approximately equal numbers of males and females were in each group. Mice received six injections of 10 µg/kg body weight (~80 U/kg) at 7 day intervals, followed by a final injection of 25 µg/kg body weight (~200 U/kg) one week after the sixth dose. Blood was collected into 1/10 volume 3.8% trisodium citrate by terminal cardiac puncture 4 days after the final injection. Samples were placed on ice before centrifugation at 3000 x g for 15 minutes at 4°C to collect plasma.

[00151] *FVIII coagulation assays and ELISAs* – FVIII activity was measured by one-stage clotting assay (Bowie and Owen, 1984, in Disorders of Hemostasis, O.D.

Ratnoff and Forbes, C.D., editors, Grune & Stratton, Inc., Orlando, 43-72) as described previously (Doering et al., 2002, *J. Biol. Chem.* 277:38345-38349) using normal human plasma (FACT) as the standard. In some assays the activated partial thromboplastin time reagent was diluted from the standard concentration. FVIII inhibitor titers were measured by a modified Bethesda assay in which fVIII constructs were added to hemophilia A plasma to a final concentration of 0.8 – 1.2 units per ml incubated with varying concentrations of inhibitor plasma for 2 hours at 37°C. One Bethesda unit (BU) is defined as the amount of inhibitory activity that produces 50% inhibition of fVIII activity in the one-stage clotting assay. The 50% inhibition point was identified by interpolation using only data points falling within a range of 40-60% inhibition. An average of at least three data points in this range were used for each determination.

[00152] Antibodies to HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 were measured by sandwich ELISA using immobilized isologous fVIII antigens. Murine monoclonal antibodies ESH4 and biotinylated-ESH8 were used as the capture and detection antibodies, respectively, as described previously (Lubin et al., 1994, *J. Biol. Chem.* 269:8639-8641). Absorbance values obtained from seven dilution of test plasma were plotted versus the logarithm of the plasma dilution and the resulting sigmoidal curves were fit to modified version of the 4-parameter logistic fit equation by nonlinear regression using the Levenberg-Marquardt algorithm. The ELISA titer was defined empirically as the dilution of plasma that returns an absorbance value of 0.3 derived from the fitted curves.

[00153] *Construction, expression and purification of recombinant proteins* – The C2 domain of fVIII contains three hydrophobic loops that comprise the binding site for phospholipid membranes (Pratt et al., (1999) *Nature* 402:439-442). Loops 1 and 2, corresponding to amino acids M2199/F2200 and L2251/L2252 in human fVIII are antigenic (Barow et al., (2001) *Blood* 97:169-174). Amino acids were selected for replacement of human residues at loops 1 and 2 based on the sequences of murine, canine, porcine, chicken and pufferfish (*Fugu rubripes*) fVIII (Table 3). The rationale for making substitutions with animal amino acids was that phospholipid membrane binding may have been preserved during evolutionary drift, thereby avoiding loss of

function due to the mutations. The amino acid replacements for the A2C2epi1, A2C2epi2 and A2C2epi3 constructs is shown in Fig. 6.

Table 3: Amino acids at phospholipid binding loops 1 and 2 in fVIII

	Amino Acid Residue			
	Loop 1		Loop 2	
Species	2199	2200	2251	2252
Human	M	F	L	L
Mouse	M	F	L	F
Dog	M	L	L	L
Pig	I	F	L	L
Chicken	I	F	V	F
Pufferfish	L	L	L	L

[00154] FVIII cDNAs were transfected into BHK-derived cells and selected for neomycin resistance. The highest expressing clones for HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 were expanded into triple flasks for production of fVIII. Cell culture supernatants from each construct were collected every 24 or 48 h for three to eleven days and pooled. The relative expression of A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 was two- to four- fold lower than HBD (Fig. 7). Two production runs of A2C2epi1 and A2C2epi3 were done to obtain sufficient material for the study.

[00155] The fVIII constructs were isolated by ion-exchange chromatography. Because the lower yields of A2C2epi1, A2C2epi2 and A2C2epi3 equate to a larger amount of impurities in the starting material, an additional mono S ion-exchange chromatography step was added. Purification tables for the constructs are shown in Table 4. The yields of the different constructs were similar through the Source Q chromatography step. The overall yields of A2C2epi1 and A2C2epi3 were lower because of the additional mono S chromatography step. The activation quotients (AQs) for the preparations ranged from 11 to 23. The AQ of A2C2epi1 and A2C2epi3 were somewhat lower than HBD, the significance of which is not known. The specific coagulant activity (U/A₂₈₀) of A2C2epi1 was lower than HBD. The specific activities of the other fVIII constructs were similar to HBD with the exception of A2C2epi2, which was approximately 25% higher.

[00156] SDS-page analysis of the preparation is shown in Fig. 8. The preparations were highly purified and appeared to be similar in composition. All contained a slight amount of a low molecular weight contaminant (marked by “*”) and a small amount of single chain fVIII. All of the preparations were cleaved by thrombin (factor IIa) to produce characteristic A1, A2 and thrombin-cleaved light chain subunits.

Table 4: Purification tables for HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3

HBD

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	12000	2.911	34927	1.50	18000	0.5	29	100%	1
SP-Sephadex pool	285	0.030	8.408	43.4	12369	1471	16	69%	2855
Source Q Pool	5.6	0.398	2.229	1287.0	7207	3234	21	40%	6275

A2epi7

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	12,600	2.802	35305	0.66	8316	0.2	9.4	100%	1
SP-Sephadex pool	114	0.060	6.840	43.0	4902	717	25	59%	3043
Source Q Pool	3.8	0.162	0.616	590.0	2242	3642	20	27%	15462

A2C2epi1 #1

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	8,400	2.802	23537	0.33	2772	0.1	7	100%	1
SP-Sephadex pool	148	0.028	4.144	4.0	592	143	16	21%	1213
Source Q Pool	10	0.081	0.810	65.00	650	802	17	23%	8025
Mono S Pool	3.2	0.109	0.349	211.0	675	1936	10	24%	19358

A2C2epi1 #2

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	10,000	2.800	28,000	0.15	1450	0.1	6	100%	1
SP-Sephadex pool	202	0.033	6.666	5.1	1030	155	11	71%	2984
Source Q Pool	4.8	0.139	0.667	105.0	504	755	9	35%	14587
Mono S Pool	3.2	0.023	0.074	72.4	232	3148	23	16%	60535

A2C2epi2

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	8,000	2.802	22,416	0.73	5840	0.3	12	100%	1
SP- Sephadex pool	162	0.019	3.013	25.2	4082	1355	17	70%	5200
Source Q Pool	10	0.102	1.020	240.00	2400	2353	24	41%	9031
Mono S Pool	3.1	0.180	0.558	790.0	2449	4389	23	42%	14630

A2C2epi3 #1

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	8,400	2.802	23537	0.27	2285	0.1	10	100%	1
SP- Sephadex pool	175	0.018	3.150	4.4	770	244	13	34%	2518
Source Q Pool	9	0.070	0.630	70.6	635	1009	10	28%	10390
Mono S Pool	3.2	0.053	0.170	152.0	486	2868	15	21%	28679

A2C2epi3 #2

Sample	Volume †	A ₂₈₀	Total A ₂₈₀	Activity ‡	Units	Units/ A ₂₈₀	AQ	% Yield	Fold Pur.
Media	12,500	2.802	35025	0.22	2800	0.1	8	100%	1
SP- Sephadex pool	148	0.041	6.068	11.0	1628	268	9	58%	3356
Source Q Pool	8.8	0.198	1.742	158.0	1390	798	12	50%	9982
Mono S Pool	3.2	0.111	0.355	283.0	906	2550	11	16%	31869

† milliliters

‡ units/milliliter

[00157] *Comparative immunogenicity A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 in hemophilia A mice* – The immunogenicity of HBD, A2epi7, A2C2epi1, A2C2epi2 and A2C2epi3 were compared in a randomized trial in hemophilia A mice. Antibodies to the immunogens were measured by Bethesda assay (Fig. 9) and by anti-fVIII ELISA (Fig. 10). The isologous form of fVIII was used as the antigen. The Bethesda titers of the A2epi7, A2C2epi1 and A2C2epi3 were significantly less than the HBD group ($p < 0.0001$, $p = 0.004$, and $p < 0.0001$, respectively, Mann-Whitney U test). In contrast, the immunogenicity of the A2C2epi2 group was not significantly lower than the HBD group. Additionally, the Bethesda titer of the A2C2epi3 group was significantly lower than the A2epi7 group ($p = 0.009$). The average Bethesda titer of the A2C2epi3 group was only 6.8, compared to average titers of 290, 62, 110 and 270 for the HBD, A2epi7, A2C2epi1 and A2C2epi2 groups, respectively.

[00158] By ELISA, the immunogenicity rank order also was HBD > A2epi7 > A2C2epi3. The results of all pair wise comparisons within this rank order were statistically significant ($p < 0.01$). However, in contrast to the Bethesda assay, the titer of A2C2epi1 in the ELISA assay was not significantly less than HBD. The immunogenicity of A2C2epi3 was profoundly depressed in the ELSA with only 3 of 25 mice testing positive.

[00159] *Antigenicity of HBD in A2C2epi3-immunized mice* – The Bethesda and ELISAs were performed using the isologous form of fVIII as the antigen. For example, in A2C2epi3-immunized mice, purified A2C2epi3 was used to reconstitute hemophilia A plasma in the Bethesda assay and was used to coat ELISA plates. To guard against the possibility that A2C2epi3 immunized mice might have developed a strong cross-reactive response to human fVIII, ten plasma samples were randomly selected from the A2C2epi3 group and tested by Bethesda assay using hemophilia A plasma reconstituted with A2C2epi3 or HBD. Table 5 shows that the immune response to HBD was less than or equal to A2C2epi3 in all of the mice. This confirms that the plasmas in the A2C2epi3 group are poorly reactive with human fVIII inhibitory epitopes.

Table 5: Bethesda titers of HBD in A2C2epi3 immunized mice

Mouse ID	ANTIGEN	
	A2C2epi3	HBD
2139-3	13	< 1.5
2134-1	13	13
2169-1	38	19
2123-2	< 1.5	< 1.5
2125-4	< 1.5	< 1.5
2122-4	< 1.5	< 1.5
2121-3	< 1.5	< 1.5
2124-2	< 1.5	< 1.5
2129-3	< 1.5	< 1.5
2141-1	< 1.5	< 1.5

[00160] *Domain specificity of anti-fVIII antibodies in hemophilia A mice immunized with HBD, A2epi7 and A2C2epi3* – Single human domain hybrid human/porcine fVIII constructs (Fig. 11) were expressed in baby hamster kidney – derived cells, purified and immobilized on microtiter plates. ELISA titers from ten plasmas from each of the

HBD and A2epi7 groups were measured. Only five plasmas in the A2C2epi3 group had significantly positive ELISA titers and were selected for study. Figure 12 shows antibodies from plasmas from the HBD and A2epi7 groups recognize all of the human domains that were studied (A1, A2, A3, C1 and C2).

[00161] There was a marked reduction in total anti-A2C2epi3 antibodies by ELISA (Figure 10). Consistent with this, there was a reduction in antibodies to the A1, A2, A3, C1 and C2 domains by domain-specific ELISA (Figure 12). If the immune system indeed is less able to respond to A2C2epi3, this indicates that anti-fVIII antibodies may develop by epitope spreading, which has been described for other immunogens (McKluskey et al., (1998) *Immunol. Rev.* 164:209-229). Thus, in the absence of initial strong recognition of the C2 domain the overall immune response may be blunted. There was also a trend toward a lower anti-A2 titer in the A2epi7 group ($p = 0.11$, Mann-Whitney U test). A hybrid human/porcine fVIII molecule that contains the A2epi7 triple mutation can be constructed in order to determine whether the sequence bounded by residues 484 – 508 is recognized less strongly in A2epi7 – immunized mice.

[00162] *In vitro recovery of fVIII activity* – The low immunogenicity of A2C2epi3 and A2epi7 could be due to denaturation of the proteins in murine hemophilia A plasma, resulting in poor delivery of the immunogen to the immune system. As a first step to test this possibility, the *in vitro* recovery of fVIII activity of the test constructs was measured. Data are available from the Bethesda assays, which were done by reconstruction of human hemophilia A plasma with the isologous constructs. After reconstitution, plasmas were assayed to determine the recovery of fVIII based in the activity of the purified material. This was done on at least eight occasions for each construct. Figure 11 shows the recovery of activity was within the expected range for all of the constructs.

[00163] *Phospholipid-dependent activated partial thromboplastin times of A2C2epi3 and HBD* – Because A2C2epi3 contains mutations at sites known to be important for the binding of fVIII to phospholipid, it is possible that it is defective with respect to this function. The specific activities of two independent preparations of A2C2epi3 were 2870 and 2550 U/A₂₈₀, compared to 3230 U/A₂₈₀ for HBD. These values are considered within experimental error, indicating that A2C2epi3 is fully

functional. Activated partial thromboplastin (aPTT)-based fVIII assays are performed at high concentrations of phospholipid. This may drive the equilibrium of forms of activated fVIII that have lower affinities for phospholipid to the bound site. To test this possibility, the aPTTs were performed on human hemophilia A plasmas reconstituted with HBD or A2C2epi3 at various dilutions of aPTT reagent, which contains phospholipid and a contact activation surface (Fig. 12). Dilutions of HBD and A2C2epi3 were made that result in equal clotting times using undiluted aPTT reagent. As the concentrations of aPTT reagent decreased below 0.5 of normal, there was an increase in clotting time for both HBD and A2C2epi3, indicating that phospholipid was a limiting component in the assay. There was a slight increase in the clotting time of A2C2epi3 compared to HBD at dilutions of aPTT reagent up to four-fold and then a more pronounced increase at higher dilutions. This result indicates that A2C2epi3 may have a lower affinity for phospholipid membranes than HBD.

[00164] While the invention has been described with certain preferred embodiments, it is understood that the preceding description is not intended to limit the scope of the invention. It will be appreciated by one skilled in the art that various equivalents and modifications can be made to the invention shown in the specific embodiments without departing from the spirit and scope of the invention. All references are incorporated herein to the extent not inconsistent.

We Claim:

1. A modified human fVIII comprising an immunoreactivity reducing amino acid substitution at each of positions number 484, 489 and 492 as set forth in SEQ ID NO: 2.
2. The modified human fVIII of claim 1 wherein each immunoreactivity reducing amino acid substitution is selected independently from the group consisting of alanine, methionine, leucine, serine, and glycine.
3. The modified human factor VIII of claim 2 wherein each immunoreactivity reducing amino acid substitution is alanine.
4. The modified human factor VIII of claim 2 wherein each immunoreactivity reducing amino acid substitution is the corresponding amino acid from porcine fVIII as set forth in SEQ ID NO: 3.
5. The modified human fVIII of claim 1 comprising a partial deletion of the B domain.
6. The modified human fVIII of claim 1 wherein the partial B-domain deletion consists of deletion of amino acids 746-1639 as set forth on SEQ ID NO: 2.
7. The modified human fVIII of claim 1 wherein said modified fVIII is a continuous single polypeptide.
8. The modified human fVIII of claim 1 wherein said modified fVIII is a A1/A2/A3-C1-C2 heterotrimer.
9. The modified human fVIII of claim 1 wherein said modified fVIII is a A1-A2/A3-C1-C2 heterodimer.

10. The modified human factor VIII of claim 1 wherein leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252 as set forth on SEQ ID NO: 2.
11. A pharmacological composition comprising the modified human fVIII of claim 1.
12. A method of managing a hemophilic patient comprising administering to said patient the pharmacological composition of claim 11.
13. DNA encoding a modified human fVIII comprising an immunoreactivity reducing amino acid substitution at each of positions number 484, 489 and 492 as set forth in SEQ ID NO: 2.
14. The DNA of claim 13 wherein each immunoreactivity reducing amino acid substitution is selected independently from the group consisting of alanine, methionine, leucine, serine, and glycine.
15. The DNA of claim 14 wherein each immunoreactivity reducing amino acid substitution is alanine.
16. The DNA of claim 14 wherein each immunoreactivity reducing amino acid substitution is the corresponding amino acid from porcine fVIII as set forth in SEQ ID NO: 3.
17. The DNA of claim 13 wherein leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252 as set forth in SEQ ID NO: 2.
18. A method of producing a modified fVIII by expressing the DNA of claim 17 in a host cell.

19. The method of claim 18 wherein said DNA is a continuous DNA sequence.
20. A modified human fVIII comprising an amino acid substitution each of positions number 2199, 2200, 2251 and 2252 as set forth in SEQ ID NO: 2, wherein leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252.
21. The modified human fVIII of claim 20 comprising a partial deletion of the B domain.
22. The modified human fVIII of claim 21 wherein the partial B-domain deletion consists of deletion of amino acids 746-1639 as set forth on SEQ ID NO: 2.
23. The modified human fVIII of claim 20 wherein said modified fVIII has reduced immunogenicity as compared to unmodified fVIII.
24. A pharmacological composition comprising the modified human fVIII of claim 20.
25. A method of managing a hemophilic patient comprising administering to said patient the pharmacological composition of claim 24.
26. DNA encoding a modified human fVIII comprising modified human fVIII wherein leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252 as set forth in SEQ ID NO: 2.
27. A method of producing a modified fVIII by expressing the DNA of claim 26 in a host cell.
28. The method of claim 27 wherein said DNA is a continuous DNA sequence.

29. A hybrid fVIII comprising: the A1, A2, A3 and C1 domains of human fVIII, having an immunoreactivity reducing amino acid substitution at each of positions number 484, 489 and 492 as set forth in SEQ ID NO: 2; and the C2 domain of a non-human fVIII.
30. The hybrid fVIII of claim 29 comprising the C2 domain of porcine fVIII as set forth in SEQ ID NO: 3.
31. The hybrid fVIII of claim 29 wherein each immunoreactivity reducing amino acid substitution is alanine.
32. DNA encoding a hybrid fVIII comprising: the A1, A2, A3 and C1 domains of human fVIII, having an immunoreactivity reducing amino acid substitution at each of positions number 484, 489 and 492 as set forth in SEQ ID NO: 2; and the C2 domain of a non-human fVIII.
33. The DNA of claim 32 wherein said hybrid fVIII comprises the C2 domain of porcine fVIII as set forth in SEQ ID NO: 3.
34. A hybrid fVIII comprising:
 - (a) the A1, A3, C1 and C2 domains of human fVIII, having an amino acid substitution each of positions number 2199, 2200, 2251 and 2252 as set forth in SEQ ID NO: 2, wherein leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252; and
 - (b) the A2 domain of a non-human fVIII.
35. The hybrid fVIII of claim 34 comprising the A2 domain of porcine fVIII as set forth in SEQ ID NO: 3.
36. DNA encoding a hybrid fVIII comprising:

- (a) the A1, A3, C1 and C2 domains of human fVIII, having an amino acid substitution each of positions number 2199, 2200, 2251 and 2252 as set forth in SEQ ID NO: 2, wherein leucine is substituted for methionine 2199, leucine is substituted for phenylalanine 2200, valine is substituted for leucine 2251, and phenylalanine is substituted for leucine 2252; and
 - (b) the A2 domain of a non-human fVIII.
37. The DNA of claim 37 wherein said hybrid fVIII comprises the A2 domain of porcine fVIII as set forth in SEQ ID NO: 3.
38. A method of identifying a modified fVIII having reduced immunogenicity or antigenicity comprising the steps:
- (a) injecting at least one dose of said modified fVIII into a first group of animal test subjects;
 - (b) injecting at least one dose of an unmodified fVIII or a fVIII with known antigenic and/or immunogenic properties into a second group of animal test subjects; and
 - (c) using a diagnostic assay to compare the inhibitory antibodies produced by said first group of animal test subjects with inhibitory antibodies produced by said second group of animal test subjects.
39. The method of claim 38 wherein said animals are mice.




FIG. 1




FIG. 2B




FIG. 2A





FIG. 3



Factor VIII Construct

FIG. 4




FIG. 5

	Loop 1	Loop 2	Loop 3
Human	MF-----V-----LL		
C2 epi1	IL-----V-----LF		
C2 epi2	IL-----V-----VF		
C2 epi3	LL-----V-----VF		

FIG. 6

7/14




FIG. 7A




FIG. 7B




FIG. 7C





FIG. 7D



Day

FIG. 7E




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14

SEQUENCE LISTING

<110> Lollar, John S

<120> Modified fVIII Having Reduced Immunogenicity Through Mutagenesis of A2 and C2 Epitopes

<130> 111-03

<150> US 60/516647

<151> 2003-10-30

<160> 4

<170> PatentIn version 3.3

<210> 1

<211> 9009

<212> DNA

<213> Homo sapiens

<400> 1
 cagtgggtaa gttccttaaa tgctctgcaa agaaattggg acttttcatt aaatcagaaa 60
 ttttactttt ttcccctcct gggagctaaa gatatttttag agaagaatta accttttgct 120
 tctccagttg aacatttgta gcaataagtc atgcaaatag agctctccac ctgcttcttt 180
 ctgtgccttt tgcgattctg ctttagtgcc accagaagat actacctggg tgcagtggaa 240
 ctgtcatggg actatatgca aagtgatctc ggtgagctgc ctgtggacgc aagatttcct 300
 cctagagtgc caaaatcttt tccattcaac acctcagtcg tgtacaaaaa gactctgttt 360
 gtagaattca cggttcacct tttcaacatc gctaagccaa ggccaccctg gatgggtctg 420
 ctaggtccta ccatccaggc tgaggtttat gatacagtg tcatcact taagaacatg 480
 gcttcccatc ctgtcagtct tcatgctgtt ggtgtatcct actggaaagc ttctgagggg 540
 gctgaatatg atgatcagac cagtcaaagg gagaaagaag atgataaagt cttccctggt 600
 ggaagccata catatgtctg gcaggctctg aaagagaatg gtccaatggc ctctgaccca 660
 ctgtgcctta cctactcata tctttctcat gtggacctgg taaaagactt gaattcaggc 720
 ctcatggag ccctactagt atgtagagaa gggagtctgg ccaaggaaaa gacacagacc 780
 ttgcacaaat ttatactact ttttgctgta tttgatgaag ggaaaagttg gcaactcagaa 840
 acaaagaact ctttgatgca ggatagggat gctgcatctg ctcgggcctg gcctaaaatg 900
 cacacagtca atggttatgt aaacaggtct ctgccaggtc tgattggatg ccacaggaaa 960
 tcagtctatt ggcattgtgat tggaatgggc accactcctg aagtgcactc aatattcctc 1020
 gaaggtcaca catttcttgt gaggaacct cgccaggcgt ccttggaaat ctgcgcaata 1080
 actttcctta ctgctcaaac actcttgatg gaccttgga agtttctact gttttgtcat 1140
 atctcttccc accaacaatga tggcatggaa gcttatgtca aagtagacag ctgtccagag 1200

gaaccccaac	tacgaatgaa	aaataatgaa	gaagcggaag	actatgatga	tgatcttact	1260
gattctgaaa	tggatgtggt	caggtttcat	gatgacaact	ctccttcctt	tatccaaatt	1320
cgctcagttg	ccaagaagca	tcctaaaact	tgggtacatt	acattgctgc	tgaagaggag	1380
gactgggact	atgctccctt	agtcctcgcc	cccgatgaca	gaagttataa	aagtcaatat	1440
ttgaacaatg	gccctcagcg	gattggtagg	aagtacaaaa	aagtccgatt	tatggcatac	1500
acagatgaaa	cctttaagac	tcgtgaagct	attcagcatg	aatcaggaat	cttgggacct	1560
ttacttttatg	gggaagttgg	agacacactg	ttgattatat	ttaagaatca	agcaagcaga	1620
ccatataaca	tctaccctca	cggaatcact	gatgtccgtc	ctttgtattc	aaggagatta	1680
ccaaaagggtg	taaaacattt	gaaggatttt	ccaattctgc	caggagaaat	attcaaatat	1740
aaatggacag	tgactgtaga	agatgggcca	actaaatcag	atcctcggtg	cctgaccgcg	1800
tattactcta	gtttcgtaa	tatggagaga	gatctagctt	caggactcat	tggccctctc	1860
ctcatctgct	acaaagaatc	tgtagatcaa	agaggaaacc	agataatgtc	agacaagagg	1920
aatgtcatcc	tgttttctgt	atctgatgag	aaccgaagct	ggtacctcac	agagaatata	1980
caacgctttc	tccccaatcc	agctggagtg	cagcttgagg	atccagagtt	ccaagcctcc	2040
aacatcatgc	acagcatcaa	tggctatggt	tttgatagtt	tgcagttgtc	agtttgtttg	2100
catgagggtg	catactggta	cattctaagc	attggagcac	agactgactt	cctttctgtc	2160
ttcttctctg	gatatacctt	caaacacaaa	atgggtctatg	aagacacact	caccctattc	2220
ccattctcag	gagaaactgt	cttcatgtcg	atggaaaacc	caggctctatg	gattctgggg	2280
tgccacaact	cagactttcg	gaacagaggc	atgaccgcct	tactgaaggt	ttctagttgt	2340
gacaagaaca	ctggtgatta	ttacgaggac	agttatgaag	atatttcagc	atacttgctg	2400
agtaaaaaca	atgccattga	accaagaagc	ttctcccaga	attcaagaca	ccctagcact	2460
aggcaaaagc	aatttaatgc	caccacaatt	ccagaaaatg	acatagagaa	gactgaccct	2520
tggtttgcac	acagaacacc	tatgcctaaa	atacaaaatg	tctcctctag	tgattttgtt	2580
atgctcttgc	gacagagtcc	tactccacat	gggctatcct	tatctgatct	ccaagaagcc	2640
aaatgatgaga	ctttttctga	tgatccatca	cctggagcaa	tagacagtaa	taacagcctg	2700
tctgaaatga	cacacttcag	gccacagctc	catcacagtg	gggacatggg	atttaccctt	2760
gagtcaggcc	tccaattaag	attaaatgag	aaactgggga	caactgcagc	aacagagttg	2820
aagaaacttg	atttcaaagt	ttctagtaca	tcaaataatc	tgatttcaac	aattccatca	2880
gacaatttgg	cagcaggtac	tgataataca	agttccttag	gacccccaag	tatgccagtt	2940
cattatgata	gtcaattaga	taccactcta	tttggcaaaa	agtcactctc	ccttactgag	3000

tctggtagac	ctctgagctt	gagtgaagaa	aataatgatt	caaagttggt	agaatcaggt	3060
ttaatgaata	gccaagaaag	ttcatgggga	aaaaatgtat	cgtcaacaga	gagtggtagg	3120
ttattttaaag	ggaaaagagc	tcattggacct	gctttgttga	ctaaagataa	tgccttatct	3180
aaagtttagca	tctctttggt	aaagacaaac	aaaacttcca	ataattcagc	aactaataga	3240
aagactcaca	ttgatggccc	atcattatta	attgagaata	gtccatcagt	ctggcaaaat	3300
atattagaaa	gtgacactga	gtttaaaaaa	gtgacacctt	tgattcatga	cagaatgctt	3360
atggacaaaa	atgctacagc	tttgaggcta	aatcatatgt	caaataaaac	tacttcatca	3420
aaaaacatgg	aaatgggtcca	acagaaaaaa	gagggcccca	ttccaccaga	tgcacaaaat	3480
ccagatatgt	cgttctttta	gatgctatct	ttgccagaat	cagcaagggtg	gatacaaagg	3540
actcatggaa	agaactctct	gaactctggg	caaggcccca	gtccaaagca	attagtatcc	3600
ttaggaccag	aaaaatctgt	ggaagggtcag	aatttcttgt	ctgagaaaaa	caaagtggtg	3660
gtaggaaagg	gtgaatttac	aaaggacgta	ggactcaaag	agatgggtttt	tccaagcagc	3720
agaaacctat	ttcttactaa	cttggataat	ttacatgaaa	ataatacaca	caatcaagaa	3780
aaaaaaattc	aggaagaaat	agaaaagaag	gaaacattaa	tccaagagaa	tgtagtgtttg	3840
cctcagatac	atacagtga	tggcactaag	aatttcatga	agaacctttt	cttactgagc	3900
actaggcaaa	atgtagaagg	ttcatatgag	ggggcatatg	ctccagtact	tcaagatttt	3960
agggtcattaa	atgattcaac	aaatagaaca	aagaaacaca	cagctcattt	ctcaaaaaaa	4020
ggggaggaag	aaaacttgga	aggcttggga	aatcaaacca	agcaaattgt	agagaaatat	4080
gcatgcacca	caaggatata	tcctaataca	agccagcaga	attttgtcac	gcaacgtagt	4140
aagagagctt	tgaacaatt	cagactccca	ctagaagaaa	cagaacttga	aaaaaggata	4200
attgtggatg	acacctcaac	ccagtgggtc	aaaaacatga	aacatttgac	cccgagcacc	4260
ctcacacaga	tagactacaa	tgagaaggag	aaagggggcca	ttactcagtc	tcccttatca	4320
gattgcctta	cgaggagtca	tagcatccct	caagcaaata	gatctccatt	accatttgca	4380
aaggatatcat	catttccatc	tattagacct	atatatctga	ccagggctct	attccaagac	4440
aactcttctc	atcttccagc	agcatcttat	agaaagaaag	attctggggg	ccaagaaagc	4500
agtcatttct	tacaaggagc	caaaaaaaat	aacctttctt	tagccattct	aaccttggag	4560
atgactgggtg	atcaaagaga	ggttgggtcc	ctgggggacaa	gtgccacaaa	ttcagtcaca	4620
tacaagaaag	ttgagaacac	tgttctcccg	aaaccagact	tgcccaaaac	atctggcaaa	4680
gttgaattgc	ttccaaaagt	tcacatttat	cagaaggacc	tattccctac	ggaaactagc	4740
aatgggtctc	ctggccatct	ggatctctgt	gaagggagcc	ttcttcaggg	aacagagggg	4800
gcgattaagt	ggaatgaagc	aaacagacct	ggaaaagtcc	cctttctgag	agtagcaaca	4860

gaaagctctg	caaagactcc	ctccaagcta	ttggatcctc	ttgcttggga	taaccactat	4920
ggtactcaga	taccaaaaga	agagtggaaa	tcccaagaga	agtcaccaga	aaaaacagct	4980
tttaagaaaa	aggataccat	tttgtccctg	aacgcttgtg	aaagcaatca	tgcaatagca	5040
gcaataaatg	agggacaaaa	taagcccga	atagaagtca	cctgggcaaa	gcaaggtagg	5100
actgaaaggc	tgtgctctca	aaaccaccca	gtcttgaaac	gccatcaacg	ggaaataact	5160
cgtactactc	ttcagtcaga	tcaagaggaa	attgactatg	atgataccat	atcagttgaa	5220
atgaagaagg	aagatthttga	catttatgat	gaggatgaaa	atcagagccc	ccgcagcttt	5280
caaaagaaaa	cacgacacta	ttttattgct	gcagtggaga	ggctctggga	ttatgggatg	5340
agtagctccc	cacatgttct	aagaaacagg	gctcagagtg	gcagtgtccc	tcagttcaag	5400
aaagttgttt	tccaggaatt	tactgatggc	tcctttactc	agcccttata	ccgtggagaa	5460
ctaaatgaac	atthgggact	cctggggcca	tatataagag	cagaagttga	agataatatc	5520
atggtaactt	tcagaaatca	ggcctctcgt	ccctattcct	tctattctag	ccttatttct	5580
tatgaggaag	atcagaggca	aggagcagaa	cctagaaaaa	actttgtcaa	gcctaataag	5640
acaaaaactt	actthttggaa	agtgaacat	catatggcac	ccactaaaga	tgagthtgac	5700
tgcaaagcct	gggcttattt	ctctgatgtt	gacctggaaa	aagatgtgca	ctcaggcctg	5760
attggacccc	ttctgggtctg	ccacactaac	acactgaacc	ctgctcatgg	gagacaagtg	5820
acagtacagg	aatttgctct	gtttttcacc	atctthgatg	agacccaaaag	ctggtaactt	5880
actgaaaata	tggaaagaaa	ctgcagggtc	ccctgcaata	tccagatgga	agatccact	5940
tttaaagaga	attatcgctt	ccatgcaatc	aatggctaca	taatggatac	actacctggc	6000
ttagtaatgg	ctcaggatca	aaggattcga	tggtatctgc	tcagcatggg	cagcaatgaa	6060
aacatccatt	ctattcattt	cagtggacat	gtgttcactg	tacgaaaaaa	agaggagtat	6120
aaaatggcac	tgtacaatct	ctatccaggt	gtttttgaga	cagtggaaat	gttaccatcc	6180
aaagctggaa	tttggcgggt	ggaatgcctt	attggcgagc	atctacatgc	tgggatgagc	6240
acactthttc	tgggtgtacag	caataagtgt	cagactcccc	tgggaatggc	ttctggacac	6300
attagagatt	ttcagattac	agcttcagga	caatatggac	agtgggcccc	aaagctggcc	6360
agacttcatt	attccggatc	aatcaatgcc	tggagcacca	aggagccctt	ttcttgatc	6420
aaggtggatc	tgthggcacc	aatgattatt	cacggcatca	agaccaggg	tgcccgtcag	6480
aagthtcca	gcctctacat	ctctcagtht	atcatcatgt	atagthttga	tgggaagaag	6540
tggcagactt	atcgaggaaa	ttccactgga	accttaatgg	tcttctthtg	caatgtggat	6600
tcactctggga	taaaacacaa	tattthtaac	cctccaatta	ttgctcgata	catccgthtg	6660

cacccaactc attatagcat tcgcagcact cttcgcatgg agttgatggg ctgtgattta	6720
aatagttgca gcatgccatt gggaatggag agtaaagcaa tatcagatgc acagattact	6780
gcttcacccct actttaccaa tatgtttgcc acctgggtctc cttcaaaagc tcgacttcac	6840
ctccaaggga ggagtaatgc ctggagacct caggtgaata atccaaaaga gtggctgcaa	6900
gtggacttcc agaagacaat gaaagtcaca ggagtaacta ctcagggagt aaaatctctg	6960
cttaccagca tgtatgtgaa ggagttcctc atctccagca gtcaagatgg ccatcagtgg	7020
actctctttt ttcagaatgg caaagtaaag gtttttcagg gaaatcaaga ctccctcaca	7080
cctgtggtga actctctaga cccaccgtta ctgactcgct accttcgaat tcacccccag	7140
agttgggtgc accagattgc cctgaggatg gaggttctgg gctgcgaggc acaggacctc	7200
tactgagggg ggccactgca gcacctgcca ctgccgtcac ctctccctcc tcagctccag	7260
ggcagtgtcc ctccctggct tgccttctac ctttgtgcta aatcctagca gacactgcct	7320
tgaagcctcc tgaattaact atcatcagtc ctgcatttct ttgggtggggg gccaggaggg	7380
tgcattccaat ttaacttaac tcttacctat tttctgcagc tgctcccaga ttactccttc	7440
cttccaatat aactaggcaa aaagaagtga ggagaaacct gcatgaaagc attcttccct	7500
gaaaagttag gcctctcaga gtcaccactt cctctgttgt agaaaaacta tgtgatgaaa	7560
ctttgaaaaa gatatttatg atgttaacat ttcagggtta gcctcatagc tttaaaataa	7620
aactctcagt tgtttattat cctgatcaag catggaacaa agcatgtttc aggatcagat	7680
caatacaatc ttggagtcaa aaggcaaatc atttggacaa tctgcaaaat ggagagaata	7740
caataactac tacagtaaag tctgtttctg cttccttaca catagatata attatgttat	7800
ttagtcatta tgagggggcac attcttatct ccaaaactag cattcttaaa ctgagaatta	7860
tagatggggg tcaagaatcc ctaagtcccc tgaaattata taaggcattc tgtataaatg	7920
caaatgtgca tttttctgac gagtgtccat agatataaag ccattgggtct taattctgac	7980
caataaaaaa ataagtcagg aggatgcaat tgttgaaagc tttgaaataa aataacatgt	8040
cttcttgaaa tttgtgatgg ccaagaaaga aaatgatgat gacattaggc ttctaaagga	8100
catacattta atatttctgt ggaaatatga ggaaaatcca tggttatctg agataggaga	8160
tacaaacttt gtaattctaa taatgcactc agtttactct ctccctctac taatttctctg	8220
ctgaaaataa cacaacaaaa atgtaacagg ggaaattata taccgtgact gaaaactaga	8280
gtcctactta catagttgaa atatcaagga ggtcagaaga aaattggact ggtgaaaaca	8340
gaaaaaacac tccagtctgc catatcacca cacaatagga tcccccttct tgccctccac	8400
cccataaga ttgtgaaggg tttactgctc cttccatctg cctgcacccc ttcactatga	8460
ctacacagaa ctctcctgat agtaaagggg gctggaggca aggataagtt atagagcagt	8520

```

tggaggaagc atccaaagac tgcaaccacag ggcaaattgga aaacaggaga tcctaatatg      8580
aaagaaaaat ggatcccaat ctgagaaaag gcaaaagaat ggctactttt ttctatgctg      8640
gagtattttc taataatcct gcttgaccct tatctgacct ctttggaaac tataacatag      8700
ctgtcacagt atagtcacaa tccacaaatg atgcagggtgc aaatgggttta tagccctgtg      8760
aagttcttaa agtttagagg ctaacttaca gaaatgaata agttgttttg ttttatagcc      8820
cggtagagga gttaacccca aaggatgat ggttttatit cctgttatgt ttaacttgat      8880
aatcttattt tggcattctt ttccattga ctatatacat ctctatttct caaatgttca      8940
tggaaactagc tcttttattt tctgtctggt ttcttcagta atgagttaaa taaaacattg      9000
acacataca                                     9009

```

```

<210> 2
<211> 2332
<212> PRT
<213> Homo sapiens

```

```

<400> 2

```

```

Ala Thr Arg Arg Tyr Tyr Leu Gly Ala Val Glu Leu Ser Trp Asp Tyr
1           5           10           15

```

```

Met Gln Ser Asp Leu Gly Glu Leu Pro Val Asp Ala Arg Phe Pro Pro
          20           25           30

```

```

Arg Val Pro Lys Ser Phe Pro Phe Asn Thr Ser Val Val Tyr Lys Lys
35           40           45

```

```

Thr Leu Phe Val Glu Phe Thr Val His Leu Phe Asn Ile Ala Lys Pro
50           55           60

```

```

Arg Pro Pro Trp Met Gly Leu Leu Gly Pro Thr Ile Gln Ala Glu Val
65           70           75           80

```

```

Tyr Asp Thr Val Val Ile Thr Leu Lys Asn Met Ala Ser His Pro Val
          85           90           95

```

```

Ser Leu His Ala Val Gly Val Ser Tyr Trp Lys Ala Ser Glu Gly Ala
100           105           110

```

```

Glu Tyr Asp Asp Gln Thr Ser Gln Arg Glu Lys Glu Asp Asp Lys Val
115           120           125

```

```

Phe Pro Gly Gly Ser His Thr Tyr Val Trp Gln Val Leu Lys Glu Asn
130           135           140

```

Gly Pro Met Ala Ser Asp Pro Leu Cys Leu Thr Tyr Ser Tyr Leu Ser
 145 150 155 160

His Val Asp Leu Val Lys Asp Leu Asn Ser Gly Leu Ile Gly Ala Leu
 165 170 175

Leu Val Cys Arg Glu Gly Ser Leu Ala Lys Glu Lys Thr Gln Thr Leu
 180 185 190

His Lys Phe Ile Leu Leu Phe Ala Val Phe Asp Glu Gly Lys Ser Trp
 195 200 205

His Ser Glu Thr Lys Asn Ser Leu Met Gln Asp Arg Asp Ala Ala Ser
 210 215 220

Ala Arg Ala Trp Pro Lys Met His Thr Val Asn Gly Tyr Val Asn Arg
 225 230 235 240

Ser Leu Pro Gly Leu Ile Gly Cys His Arg Lys Ser Val Tyr Trp His
 245 250 255

Val Ile Gly Met Gly Thr Thr Pro Glu Val His Ser Ile Phe Leu Glu
 260 265 270

Gly His Thr Phe Leu Val Arg Asn His Arg Gln Ala Ser Leu Glu Ile
 275 280 285

Ser Pro Ile Thr Phe Leu Thr Ala Gln Thr Leu Leu Met Asp Leu Gly
 290 295 300

Gln Phe Leu Leu Phe Cys His Ile Ser Ser His Gln His Asp Gly Met
 305 310 315 320

Glu Ala Tyr Val Lys Val Asp Ser Cys Pro Glu Glu Pro Gln Leu Arg
 325 330 335

Met Lys Asn Asn Glu Glu Ala Glu Asp Tyr Asp Asp Asp Leu Thr Asp
 340 345 350

Ser Glu Met Asp Val Val Arg Phe Asp Asp Asp Asn Ser Pro Ser Phe
 355 360 365

Ile Gln Ile Arg Ser Val Ala Lys Lys His Pro Lys Thr Trp Val His
 370 375 380

Tyr Ile Ala Ala Glu Glu Glu Asp Trp Asp Tyr Ala Pro Leu Val Leu
 385 390 395 400

Ala Pro Asp Asp Arg Ser Tyr Lys Ser Gln Tyr Leu Asn Asn Gly Pro
 405 410 415

Gln Arg Ile Gly Arg Lys Tyr Lys Lys Val Arg Phe Met Ala Tyr Thr
 420 425 430

Asp Glu Thr Phe Lys Thr Arg Glu Ala Ile Gln His Glu Ser Gly Ile
 435 440 445

Leu Gly Pro Leu Leu Tyr Gly Glu Val Gly Asp Thr Leu Leu Ile Ile
 450 455 460

Phe Lys Asn Gln Ala Ser Arg Pro Tyr Asn Ile Tyr Pro His Gly Ile
 465 470 475 480

Thr Asp Val Arg Pro Leu Tyr Ser Arg Arg Leu Pro Lys Gly Val Lys
 485 490 495

His Leu Lys Asp Phe Pro Ile Leu Pro Gly Glu Ile Phe Lys Tyr Lys
 500 505 510

Trp Thr Val Thr Val Glu Asp Gly Pro Thr Lys Ser Asp Pro Arg Cys
 515 520 525

Leu Thr Arg Tyr Tyr Ser Ser Phe Val Asn Met Glu Arg Asp Leu Ala
 530 535 540

Ser Gly Leu Ile Gly Pro Leu Leu Ile Cys Tyr Lys Glu Ser Val Asp
 545 550 555 560

Gln Arg Gly Asn Gln Ile Met Ser Asp Lys Arg Asn Val Ile Leu Phe
 565 570 575

Ser Val Phe Asp Glu Asn Arg Ser Trp Tyr Leu Thr Glu Asn Ile Gln
 580 585 590

Arg Phe Leu Pro Asn Pro Ala Gly Val Gln Leu Glu Asp Pro Glu Phe
 595 600 605

Gln Ala Ser Asn Ile Met His Ser Ile Asn Gly Tyr Val Phe Asp Ser
 610 615 620

Leu Gln Leu Ser Val Cys Leu His Glu Val Ala Tyr Trp Tyr Ile Leu
 625 630 635 640

Ser Ile Gly Ala Gln Thr Asp Phe Leu Ser Val Phe Phe Ser Gly Tyr
 645 650 655

Thr Phe Lys His Lys Met Val Tyr Glu Asp Thr Leu Thr Leu Phe Pro
 660 665 670

Phe Ser Gly Glu Thr Val Phe Met Ser Met Glu Asn Pro Gly Leu Trp
 675 680 685

Ile Leu Gly Cys His Asn Ser Asp Phe Arg Asn Arg Gly Met Thr Ala
 690 695 700

Leu Leu Lys Val Ser Ser Cys Asp Lys Asn Thr Gly Asp Tyr Tyr Glu
 705 710 715 720

Asp Ser Tyr Glu Asp Ile Ser Ala Tyr Leu Leu Ser Lys Asn Asn Ala
 725 730 735

Ile Glu Pro Arg Ser Phe Ser Gln Asn Ser Arg His Pro Ser Thr Arg
 740 745 750

Gln Lys Gln Phe Asn Ala Thr Thr Ile Pro Glu Asn Asp Ile Glu Lys
 755 760 765

Thr Asp Pro Trp Phe Ala His Arg Thr Pro Met Pro Lys Ile Gln Asn
 770 775 780

Val Ser Ser Ser Asp Leu Leu Met Leu Leu Arg Gln Ser Pro Thr Pro
 785 790 795 800

His Gly Leu Ser Leu Ser Asp Leu Gln Glu Ala Lys Tyr Glu Thr Phe
 805 810 815

Ser Asp Asp Pro Ser Pro Gly Ala Ile Asp Ser Asn Asn Ser Leu Ser
 820 825 830

Glu Met Thr His Phe Arg Pro Gln Leu His His Ser Gly Asp Met Val
 835 840 845

Phe Thr Pro Glu Ser Gly Leu Gln Leu Arg Leu Asn Glu Lys Leu Gly
 850 855 860

Thr Thr Ala Ala Thr Glu Leu Lys Lys Leu Asp Phe Lys Val Ser Ser

865		870		875		880
Thr Ser Asn Asn Leu Ile Ser Thr Ile Pro Ser Asp Asn Leu Ala Ala						
		885		890		895
Gly Thr Asp Asn Thr Ser Ser Leu Gly Pro Pro Ser Met Pro Val His						
		900		905		910
Tyr Asp Ser Gln Leu Asp Thr Thr Leu Phe Gly Lys Lys Ser Ser Pro						
		915		920		925
Leu Thr Glu Ser Gly Gly Pro Leu Ser Leu Ser Glu Glu Asn Asn Asp						
		930		935		940
Ser Lys Leu Leu Glu Ser Gly Leu Met Asn Ser Gln Glu Ser Ser Trp						
		945		950		955
Gly Lys Asn Val Ser Ser Thr Glu Ser Gly Arg Leu Phe Lys Gly Lys						
		965		970		975
Arg Ala His Gly Pro Ala Leu Leu Thr Lys Asp Asn Ala Leu Phe Lys						
		980		985		990
Val Ser Ile Ser Leu Leu Lys Thr Asn Lys Thr Ser Asn Asn Ser Ala						
		995		1000		1005
Thr Asn Arg Lys Thr His Ile Asp Gly Pro Ser Leu Leu Ile Glu						
		1010		1015		1020
Asn Ser Pro Ser Val Trp Gln Asn Ile Leu Glu Ser Asp Thr Glu						
		1025		1030		1035
Phe Lys Lys Val Thr Pro Leu Ile His Asp Arg Met Leu Met Asp						
		1040		1045		1050
Lys Asn Ala Thr Ala Leu Arg Leu Asn His Met Ser Asn Lys Thr						
		1055		1060		1065
Thr Ser Ser Lys Asn Met Glu Met Val Gln Gln Lys Lys Glu Gly						
		1070		1075		1080
Pro Ile Pro Pro Asp Ala Gln Asn Pro Asp Met Ser Phe Phe Lys						
		1085		1090		1095
Met Leu Phe Leu Pro Glu Ser Ala Arg Trp Ile Gln Arg Thr His						
		1100		1105		1110

Gly Lys	Asn Ser Leu Asn Ser	Gly Gln Gly Pro Ser	Pro Lys Gln
1115	1120	1125	
Leu Val	Ser Leu Gly Pro Glu	Lys Ser Val Glu Gly	Gln Asn Phe
1130	1135	1140	
Leu Ser	Glu Lys Asn Lys Val	Val Val Gly Lys Gly	Glu Phe Thr
1145	1150	1155	
Lys Asp	Val Gly Leu Lys Glu	Met Val Phe Pro Ser	Ser Arg Asn
1160	1165	1170	
Leu Phe	Leu Thr Asn Leu Asp	Asn Leu His Glu Asn	Asn Thr His
1175	1180	1185	
Asn Gln	Glu Lys Lys Ile Gln	Glu Glu Ile Glu Lys	Lys Glu Thr
1190	1195	1200	
Leu Ile	Gln Glu Asn Val Val	Leu Pro Gln Ile His	Thr Val Thr
1205	1210	1215	
Gly Thr	Lys Asn Phe Met Lys	Asn Leu Phe Leu Leu	Ser Thr Arg
1220	1225	1230	
Gln Asn	Val Glu Gly Ser Tyr	Glu Gly Ala Tyr Ala	Pro Val Leu
1235	1240	1245	
Gln Asp	Phe Arg Ser Leu Asn	Asp Ser Thr Asn Arg	Thr Lys Lys
1250	1255	1260	
His Thr	Ala His Phe Ser Lys	Lys Gly Glu Glu Glu	Asn Leu Glu
1265	1270	1275	
Gly Leu	Gly Asn Gln Thr Lys	Gln Ile Val Glu Lys	Tyr Ala Cys
1280	1285	1290	
Thr Thr	Arg Ile Ser Pro Asn	Thr Ser Gln Gln Asn	Phe Val Thr
1295	1300	1305	
Gln Arg	Ser Lys Arg Ala Leu	Lys Gln Phe Arg Leu	Pro Leu Glu
1310	1315	1320	
Glu Thr	Glu Leu Glu Lys Arg	Ile Ile Val Asp Asp	Thr Ser Thr
1325	1330	1335	

Gln Trp	Ser Lys Asn Met	Lys	His Leu Thr	Pro Ser	Thr Leu Thr
1340		1345		1350	
Gln Ile	Asp Tyr Asn Glu	Lys	Glu Lys Gly Ala	Ile	Thr Gln Ser
1355		1360		1365	
Pro Leu	Ser Asp Cys Leu	Thr	Arg Ser His Ser	Ile	Pro Gln Ala
1370		1375		1380	
Asn Arg	Ser Pro Leu Pro	Ile	Ala Lys Val Ser	Ser	Phe Pro Ser
1385		1390		1395	
Ile Arg	Pro Ile Tyr Leu	Thr	Arg Val Leu Phe	Gln	Asp Asn Ser
1400		1405		1410	
Ser His	Leu Pro Ala Ala	Ser	Tyr Arg Lys Lys	Asp	Ser Gly Val
1415		1420		1425	
Gln Glu	Ser Ser His Phe	Leu	Gln Gly Ala Lys	Lys	Asn Asn Leu
1430		1435		1440	
Ser Leu	Ala Ile Leu Thr	Leu	Glu Met Thr Gly	Asp	Gln Arg Glu
1445		1450		1455	
Val Gly	Ser Leu Gly Thr	Ser	Ala Thr Asn Ser	Val	Thr Tyr Lys
1460		1465		1470	
Lys Val	Glu Asn Thr Val	Leu	Pro Lys Pro Asp	Leu	Pro Lys Thr
1475		1480		1485	
Ser Gly	Lys Val Glu Leu	Leu	Pro Lys Val His	Ile	Tyr Gln Lys
1490		1495		1500	
Asp Leu	Phe Pro Thr Glu	Thr	Ser Asn Gly Ser	Pro	Gly His Leu
1505		1510		1515	
Asp Leu	Val Glu Gly Ser	Leu	Leu Gln Gly Thr	Glu	Gly Ala Ile
1520		1525		1530	
Lys Trp	Asn Glu Ala Asn	Arg	Pro Gly Lys Val	Pro	Phe Leu Arg
1535		1540		1545	
Val Ala	Thr Glu Ser Ser	Ala	Lys Thr Pro Ser	Lys	Leu Leu Asp
1550		1555		1560	

Pro	Leu	Ala	Trp	Asp	Asn	His	Tyr	Gly	Thr	Gln	Ile	Pro	Lys	Glu
	1565					1570					1575			
Glu	Trp	Lys	Ser	Gln	Glu	Lys	Ser	Pro	Glu	Lys	Thr	Ala	Phe	Lys
	1580					1585					1590			
Lys	Lys	Asp	Thr	Ile	Leu	Ser	Leu	Asn	Ala	Cys	Glu	Ser	Asn	His
	1595					1600					1605			
Ala	Ile	Ala	Ala	Ile	Asn	Glu	Gly	Gln	Asn	Lys	Pro	Glu	Ile	Glu
	1610					1615					1620			
Val	Thr	Trp	Ala	Lys	Gln	Gly	Arg	Thr	Glu	Arg	Leu	Cys	Ser	Gln
	1625					1630					1635			
Asn	Pro	Pro	Val	Leu	Lys	Arg	His	Gln	Arg	Glu	Ile	Thr	Arg	Thr
	1640					1645					1650			
Thr	Leu	Gln	Ser	Asp	Gln	Glu	Glu	Ile	Asp	Tyr	Asp	Asp	Thr	Ile
	1655					1660					1665			
Ser	Val	Glu	Met	Lys	Lys	Glu	Asp	Phe	Asp	Ile	Tyr	Asp	Glu	Asp
	1670					1675					1680			
Glu	Asn	Gln	Ser	Pro	Arg	Ser	Phe	Gln	Lys	Lys	Thr	Arg	His	Tyr
	1685					1690					1695			
Phe	Ile	Ala	Ala	Val	Glu	Arg	Leu	Trp	Asp	Tyr	Gly	Met	Ser	Ser
	1700					1705					1710			
Ser	Pro	His	Val	Leu	Arg	Asn	Arg	Ala	Gln	Ser	Gly	Ser	Val	Pro
	1715					1720					1725			
Gln	Phe	Lys	Lys	Val	Val	Phe	Gln	Glu	Phe	Thr	Asp	Gly	Ser	Phe
	1730					1735					1740			
Thr	Gln	Pro	Leu	Tyr	Arg	Gly	Glu	Leu	Asn	Glu	His	Leu	Gly	Leu
	1745					1750					1755			
Leu	Gly	Pro	Tyr	Ile	Arg	Ala	Glu	Val	Glu	Asp	Asn	Ile	Met	Val
	1760					1765					1770			
Thr	Phe	Arg	Asn	Gln	Ala	Ser	Arg	Pro	Tyr	Ser	Phe	Tyr	Ser	Ser
	1775					1780					1785			
Leu	Ile	Ser	Tyr	Glu	Glu	Asp	Gln	Arg	Gln	Gly	Ala	Glu	Pro	Arg

1790		1795		1800
Lys Asn Phe Val Lys Pro Asn Glu Thr Lys Thr Tyr Phe Trp Lys				
1805		1810		1815
Val Gln His His Met Ala Pro Thr Lys Asp Glu Phe Asp Cys Lys				
1820		1825		1830
Ala Trp Ala Tyr Phe Ser Asp Val Asp Leu Glu Lys Asp Val His				
1835		1840		1845
Ser Gly Leu Ile Gly Pro Leu Leu Val Cys His Thr Asn Thr Leu				
1850		1855		1860
Asn Pro Ala His Gly Arg Gln Val Thr Val Gln Glu Phe Ala Leu				
1865		1870		1875
Phe Phe Thr Ile Phe Asp Glu Thr Lys Ser Trp Tyr Phe Thr Glu				
1880		1885		1890
Asn Met Glu Arg Asn Cys Arg Ala Pro Cys Asn Ile Gln Met Glu				
1895		1900		1905
Asp Pro Thr Phe Lys Glu Asn Tyr Arg Phe His Ala Ile Asn Gly				
1910		1915		1920
Tyr Ile Met Asp Thr Leu Pro Gly Leu Val Met Ala Gln Asp Gln				
1925		1930		1935
Arg Ile Arg Trp Tyr Leu Leu Ser Met Gly Ser Asn Glu Asn Ile				
1940		1945		1950
His Ser Ile His Phe Ser Gly His Val Phe Thr Val Arg Lys Lys				
1955		1960		1965
Glu Glu Tyr Lys Met Ala Leu Tyr Asn Leu Tyr Pro Gly Val Phe				
1970		1975		1980
Glu Thr Val Glu Met Leu Pro Ser Lys Ala Gly Ile Trp Arg Val				
1985		1990		1995
Glu Cys Leu Ile Gly Glu His Leu His Ala Gly Met Ser Thr Leu				
2000		2005		2010
Phe Leu Val Tyr Ser Asn Lys Cys Gln Thr Pro Leu Gly Met Ala				
2015		2020		2025

Ser Gly	His Ile Arg Asp Phe	Gln Ile Thr Ala Ser	Gly Gln Tyr
2030	2035	2040	
Gly Gln	Trp Ala Pro Lys Leu	Ala Arg Leu His Tyr	Ser Gly Ser
2045	2050	2055	
Ile Asn	Ala Trp Ser Thr Lys	Glu Pro Phe Ser Trp	Ile Lys Val
2060	2065	2070	
Asp Leu	Leu Ala Pro Met Ile	Ile His Gly Ile Lys	Thr Gln Gly
2075	2080	2085	
Ala Arg	Gln Lys Phe Ser Ser	Leu Tyr Ile Ser Gln	Phe Ile Ile
2090	2095	2100	
Met Tyr	Ser Leu Asp Gly Lys	Lys Trp Gln Thr Tyr	Arg Gly Asn
2105	2110	2115	
Ser Thr	Gly Thr Leu Met Val	Phe Phe Gly Asn Val	Asp Ser Ser
2120	2125	2130	
Gly Ile	Lys His Asn Ile Phe	Asn Pro Pro Ile Ile	Ala Arg Tyr
2135	2140	2145	
Ile Arg	Leu His Pro Thr His	Tyr Ser Ile Arg Ser	Thr Leu Arg
2155	2160		2150
Met Glu	Leu Met Gly Cys Asp	Leu Asn Ser Cys Ser	Met Pro Leu
2165	2170	2175	
Gly Met	Glu Ser Lys Ala Ile	Ser Asp Ala Gln Ile	Thr Ala Ser
2180	2185	2190	
Ser Tyr	Phe Thr Asn Met Phe	Ala Thr Trp Ser Pro	Ser Lys Ala
2195	2200	2205	
Arg Leu	His Leu Gln Gly Arg	Ser Asn Ala Trp Arg	Pro Gln Val
2210	2215	2220	
Asn Asn	Pro Lys Glu Trp Leu	Gln Val Asp Phe Gln	Lys Thr Met
2225	2230	2235	
Lys Val	Thr Gly Val Thr Thr	Gln Gly Val Lys Ser	Leu Leu Thr
2240	2245	2250	

Ser Met Tyr Val Lys Glu Phe Leu Ile Ser Ser Ser Gln Asp Gly
 2255 2260 2265

His Gln Trp Thr Leu Phe Phe Gln Asn Gly Lys Val Lys Val Phe
 2270 2275 2280

Gln Gly Asn Gln Asp Ser Phe Thr Pro Val Val Asn Ser Leu Asp
 2285 2290 2295

Pro Pro Leu Leu Thr Arg Tyr Leu Arg Ile His Pro Gln Ser Trp
 2300 2305 2310

Val His Gln Ile Ala Leu Arg Met Glu Val Leu Gly Cys Glu Ala
 2315 2320 2325

Gln Asp Leu Tyr
 2330

<210> 3
 <211> 2114
 <212> PRT
 <213> Sus scrofa

<400> 3

Ala Ile Arg Arg Tyr Tyr Leu Gly Ala Val Glu Leu Ser Trp Asp Tyr
 1 5 10 15

Arg Gln Ser Glu Leu Leu Arg Glu Leu His Val Asp Thr Arg Phe Pro
 20 25 30

Ala Thr Ala Pro Gly Ala Leu Pro Leu Gly Pro Ser Val Leu Tyr Lys
 35 40 45

Lys Thr Val Phe Val Glu Phe Thr Asp Gln Leu Phe Ser Val Ala Arg
 50 55 60

Pro Arg Pro Pro Trp Met Gly Leu Leu Gly Pro Thr Ile Gln Ala Glu
 65 70 75 80

Val Tyr Asp Thr Val Val Val Thr Leu Lys Asn Met Ala Ser His Pro
 85 90 95

Val Ser Leu His Ala Val Gly Val Ser Phe Trp Lys Ser Ser Glu Gly
 100 105 110

Ala Glu Tyr Glu Asp His Thr Ser Gln Arg Glu Lys Glu Asp Asp Lys

115	120	125
Val Leu Pro Gly Lys Ser Gln Thr Tyr Val Trp Gln Val Leu Lys Glu		
130	135	140
Asn Gly Pro Thr Ala Ser Asp Pro Pro Cys Leu Thr Tyr Ser Tyr Leu		
145	150	155
Ser His Val Asp Leu Val Lys Asp Leu Asn Ser Gly Leu Ile Gly Ala		
165	170	175
Leu Leu Val Cys Arg Glu Gly Ser Leu Thr Arg Glu Arg Thr Gln Asn		
180	185	190
Leu His Glu Phe Val Leu Leu Phe Ala Val Phe Asp Glu Gly Lys Ser		
195	200	205
Trp His Ser Ala Arg Asn Asp Ser Trp Thr Arg Ala Met Asp Pro Ala		
210	215	220
Pro Ala Arg Ala Gln Pro Ala Met His Thr Val Asn Gly Tyr Val Asn		
225	230	235
Arg Ser Leu Pro Gly Leu Ile Gly Cys His Lys Lys Ser Val Tyr Trp		
245	250	255
His Val Ile Gly Met Gly Thr Ser Pro Glu Val His Ser Ile Phe Leu		
260	265	270
Glu Gly His Thr Phe Leu Val Arg His His Arg Gln Ala Ser Leu Glu		
275	280	285
Ile Ser Pro Leu Thr Phe Leu Thr Ala Gln Thr Phe Leu Met Asp Leu		
290	295	300
Gly Gln Phe Leu Leu Phe Cys His Ile Ser Ser His His His Gly Gly		
305	310	315
Met Glu Ala His Val Arg Val Glu Ser Cys Ala Glu Glu Pro Gln Leu		
325	330	335
Arg Arg Lys Ala Asp Glu Glu Glu Asp Tyr Asp Asp Asn Leu Tyr Asp		
340	345	350
Ser Asp Met Asp Val Val Arg Leu Asp Gly Asp Asp Val Ser Pro Phe		
355	360	365

Ile Gln Ile Arg Ser Val Ala Lys Lys His Pro Lys Thr Trp Val His
 370 375 380

Tyr Ile Ser Ala Glu Glu Glu Asp Trp Asp Tyr Ala Pro Ala Val Pro
 385 390 395 400

Ser Pro Ser Asp Arg Ser Tyr Lys Ser Leu Tyr Leu Asn Ser Gly Pro
 405 410 415

Gln Arg Ile Gly Arg Lys Tyr Lys Lys Ala Arg Phe Val Ala Tyr Thr
 420 425 430

Asp Val Thr Phe Lys Thr Arg Lys Ala Ile Pro Tyr Glu Ser Gly Ile
 435 440 445

Leu Gly Pro Leu Leu Tyr Gly Glu Val Gly Asp Thr Leu Leu Ile Ile
 450 455 460

Phe Lys Asn Lys Ala Ser Arg Pro Tyr Asn Ile Tyr Pro His Gly Ile
 465 470 475 480

Thr Asp Val Ser Ala Leu His Pro Gly Arg Leu Leu Lys Gly Trp Lys
 485 490 495

His Leu Lys Asp Met Pro Ile Leu Pro Gly Glu Thr Phe Lys Tyr Lys
 500 505 510

Trp Thr Val Thr Val Glu Asp Gly Pro Thr Lys Ser Asp Pro Arg Cys
 515 520 525

Leu Thr Arg Tyr Tyr Ser Ser Ser Ile Asn Leu Glu Lys Asp Leu Ala
 530 535 540

Ser Gly Leu Ile Gly Pro Leu Leu Ile Cys Tyr Lys Glu Ser Val Asp
 545 550 555 560

Gln Arg Gly Asn Gln Met Met Ser Asp Lys Arg Asn Val Ile Leu Phe
 565 570 575

Ser Val Phe Asp Glu Asn Gln Ser Trp Tyr Leu Ala Glu Asn Ile Gln
 580 585 590

Arg Phe Leu Pro Asn Pro Asp Gly Leu Gln Pro Gln Asp Pro Glu Phe
 595 600 605

Gln Ala Ser Asn Ile Met His Ser Ile Asn Gly Tyr Val Phe Asp Ser
 610 615 620

Leu Gln Leu Ser Val Cys Leu His Glu Val Ala Tyr Trp Tyr Ile Leu
 625 630 635 640

Ser Val Gly Ala Gln Thr Asp Phe Leu Ser Val Phe Phe Ser Gly Tyr
 645 650 655

Thr Phe Lys His Lys Met Val Tyr Glu Asp Thr Leu Thr Leu Phe Pro
 660 665 670

Phe Ser Gly Glu Thr Val Phe Met Ser Met Glu Asn Pro Gly Leu Trp
 675 680 685

Val Leu Gly Cys His Asn Ser Asp Leu Arg Asn Arg Gly Met Thr Ala
 690 695 700

Leu Leu Lys Val Tyr Ser Cys Asp Arg Asp Ile Gly Asp Tyr Tyr Asp
 705 710 715 720

Asn Thr Tyr Glu Asp Ile Pro Gly Phe Leu Leu Ser Gly Lys Asn Val
 725 730 735

Ile Glu Pro Arg Ser Phe Ala Gln Asn Ser Arg Pro Pro Ser Ala Ser
 740 745 750

Gln Lys Gln Phe Gln Thr Ile Thr Ser Pro Glu Asp Asp Val Glu Leu
 755 760 765

Asp Pro Gln Ser Gly Glu Arg Thr Gln Ala Leu Glu Glu Leu Ser Val
 770 775 780

Pro Ser Gly Asp Gly Ser Met Leu Leu Gly Gln Asn Pro Ala Pro His
 785 790 795 800

Gly Ser Ser Ser Ser Asp Leu Gln Glu Ala Arg Asn Glu Ala Asp Asp
 805 810 815

Tyr Leu Pro Gly Ala Arg Glu Arg Asn Thr Ala Pro Ser Ala Ala Ala
 820 825 830

Arg Leu Arg Pro Glu Leu His His Ser Ala Glu Arg Val Leu Thr Pro
 835 840 845

Glu Pro Glu Lys Glu Leu Lys Lys Leu Asp Ser Lys Met Ser Ser Ser
 850 855 860

Ser Asp Leu Leu Lys Thr Ser Pro Thr Ile Pro Ser Asp Thr Leu Ser
 865 870 875 880

Ala Glu Thr Glu Arg Thr His Ser Leu Gly Pro Pro His Pro Gln Val
 885 890 895

Asn Phe Arg Ser Gln Leu Gly Ala Ile Val Leu Gly Lys Asn Ser Ser
 900 905 910

His Phe Ile Gly Ala Gly Val Pro Leu Gly Ser Thr Glu Glu Asp His
 915 920 925

Glu Ser Ser Leu Gly Glu Asn Val Ser Pro Val Glu Ser Asp Gly Ile
 930 935 940

Phe Glu Lys Glu Arg Ala His Gly Pro Ala Ser Leu Thr Lys Asp Asp
 945 950 955 960

Val Leu Phe Lys Val Asn Ile Ser Leu Val Lys Thr Asn Lys Ala Arg
 965 970 975

Val Tyr Leu Lys Thr Asn Arg Lys Ile His Ile Asp Asp Ala Ala Leu
 980 985 990

Leu Thr Glu Asn Arg Ala Ser Ala Thr Phe Met Asp Lys Asn Thr Thr
 995 1000 1005

Ala Ser Gly Leu Asn His Val Ser Asn Trp Ile Lys Gly Pro Leu
 1010 1015 1020

Gly Lys Asn Pro Leu Ser Ser Glu Arg Gly Pro Ser Pro Glu Leu
 1025 1030 1035

Leu Thr Ser Ser Gly Ser Gly Lys Ser Val Lys Gly Gln Ser Ser
 1040 1045 1050

Gly Gln Gly Arg Ile Arg Val Ala Val Glu Glu Glu Glu Leu Ser
 1055 1060 1065

Lys Gly Lys Glu Met Met Leu Pro Asn Ser Glu Leu Thr Phe Leu
 1070 1075 1080

Thr Asn Ser Ala Asp Val Gln Gly Asn Asp Thr His Ser Gln Gly

1085		1090		1095	
Lys Lys Ser Arg Glu Glu Met Glu Arg Arg Glu Lys Leu Val Gln					
1100		1105		1110	
Glu Lys Val Asp Leu Pro Gln Val Tyr Thr Ala Thr Gly Thr Lys					
1115		1120		1125	
Asn Phe Leu Arg Asn Ile Phe His Gln Ser Thr Glu Pro Ser Val					
1130		1135		1140	
Glu Gly Phe Asp Gly Gly Ser His Ala Pro Val Pro Gln Asp Ser					
1145		1150		1155	
Arg Ser Leu Asn Asp Ser Ala Glu Arg Ala Glu Thr His Ile Ala					
1160		1165		1170	
His Phe Ser Ala Ile Arg Glu Glu Ala Pro Leu Glu Ala Pro Gly					
1175		1180		1185	
Asn Arg Thr Gly Pro Gly Pro Arg Ser Ala Val Pro Arg Arg Val					
1190		1195		1200	
Lys Gln Ser Leu Lys Gln Ile Arg Leu Pro Leu Glu Glu Ile Lys					1205
1210		1215			
Pro Glu Arg Gly Val Val Leu Asn Ala Thr Ser Thr Arg Trp Ser					
1220		1225		1230	
Glu Ser Ser Pro Ile Leu Gln Gly Ala Lys Arg Asn Asn Leu Ser					
1235		1240		1245	
Leu Pro Phe Leu Thr Leu Glu Met Ala Gly Gly Gln Gly Lys Ile					
1250		1255		1260	
Ser Ala Leu Gly Lys Ser Ala Ala Gly Pro Leu Ala Ser Gly Lys					
1265		1270		1275	
Leu Glu Lys Ala Val Leu Ser Ser Ala Gly Leu Ser Glu Ala Ser					
1280		1285		1290	
Gly Lys Ala Glu Phe Leu Pro Lys Val Arg Val His Arg Glu Asp					
1295		1300		1305	
Leu Leu Pro Gln Lys Thr Ser Asn Val Ser Cys Ala His Gly Asp					
1310		1315		1320	

Leu Gly	Gln Glu Ile Phe	Leu	Gln Lys Thr Arg	Gly	Pro Val Asn
1325		1330		1335	
Leu Asn	Lys Val Asn Arg	Pro	Gly Arg Thr Pro	Ser	Lys Leu Leu
1340		1345		1350	
Gly Pro	Pro Met Pro Lys	Glu	Trp Glu Ser Leu	Glu	Lys Ser Pro
1355		1360		1365	
Lys Ser	Thr Ala Leu Arg	Thr	Lys Asp Ile Ile	Ser	Leu Pro Leu
1370		1375		1380	
Asp Arg	His Glu Ser Asn	His	Ser Ile Ala Ala	Lys	Asn Glu Gly
1385		1390		1395	
Gln Ala	Glu Thr Gln Arg	Glu	Ala Ala Trp Thr	Lys	Gln Gly Gly
1400		1405		1410	
Pro Gly	Arg Leu Cys Ala	Pro	Lys Pro Pro Val	Leu	Arg Arg His
1415		1420		1425	
Gln Arg	Asp Ile Ser Leu	Pro	Thr Phe Gln Pro	Glu	Glu Asp Lys
1430		1435		1440	
Met Asp	Tyr Asp Asp Ile	Phe	Ser Thr Glu Thr	Lys	Gly Glu Asp
1445		1450		1455	
Phe Asp	Ile Tyr Gly Glu	Asp	Glu Asn Gln Asp	Pro	Arg Ser Phe
1460		1465		1470	
Gln Lys	Arg Thr Arg His	Tyr	Phe Ile Ala Ala	Val	Glu Gln Leu
1475		1480		1485	
Trp Asp	Tyr Gly Met Ser	Glu	Ser Pro Arg Ala	Leu	Arg Asn Arg
1490		1495		1500	
Ala Gln	Asn Gly Glu Val	Pro	Arg Phe Lys Lys	Val	Val Phe Arg
1505		1510		1515	
Glu Phe	Ala Asp Gly Ser	Phe	Thr Gln Pro Ser	Tyr	Arg Gly Glu
1520		1525		1530	
Leu Asn	Lys His Leu Gly	Leu	Leu Gly Pro Tyr	Ile	Arg Ala Glu
1535		1540		1545	

Val	Glu	Asp	Asn	Ile	Met	Val	Thr	Phe	Lys	Asn	Gln	Ala	Ser	Arg
1550						1555					1560			
Pro	Tyr	Ser	Phe	Tyr	Ser	Ser	Leu	Ile	Ser	Tyr	Pro	Asp	Asp	Gln
1565						1570					1575			
Glu	Gln	Gly	Ala	Glu	Pro	Arg	His	Asn	Phe	Val	Gln	Pro	Asn	Glu
1580						1585					1590			
Thr	Arg	Thr	Tyr	Phe	Trp	Lys	Val	Gln	His	His	Met	Ala	Pro	Thr
1595						1600					1605			
Glu	Asp	Glu	Phe	Asp	Cys	Lys	Ala	Trp	Ala	Tyr	Phe	Ser	Asp	Val
1610						1615					1620			
Asp	Leu	Glu	Lys	Asp	Val	His	Ser	Gly	Leu	Ile	Gly	Pro	Leu	Leu
1625						1630					1635			
Ile	Cys	Arg	Ala	Asn	Thr	Leu	Asn	Ala	Ala	His	Gly	Arg	Gln	Val
1640						1645					1650			
Thr	Val	Gln	Glu	Phe	Ala	Leu	Phe	Phe	Thr	Ile	Phe	Asp	Glu	Thr
1655						1660					1665			
Lys	Ser	Trp	Tyr	Phe	Thr	Glu	Asn	Val	Glu	Arg	Asn	Cys	Arg	Ala
1670						1675					1680			
Pro	Cys	His	Leu	Gln	Met	Glu	Asp	Pro	Thr	Leu	Lys	Glu	Asn	Tyr
1685						1690					1695			
Arg	Phe	His	Ala	Ile	Asn	Gly	Tyr	Val	Met	Asp	Thr	Leu	Pro	Gly
1700						1705					1710			
Leu	Val	Met	Ala	Gln	Asn	Gln	Arg	Ile	Arg	Trp	Tyr	Leu	Leu	Ser
1715						1720					1725			
Met	Gly	Ser	Asn	Glu	Asn	Ile	His	Ser	Ile	His	Phe	Ser	Gly	His
1730						1735					1740			
Val	Phe	Ser	Val	Arg	Lys	Lys	Glu	Glu	Tyr	Lys	Met	Ala	Val	Tyr
1745						1750					1755			
Asn	Leu	Tyr	Pro	Gly	Val	Phe	Glu	Thr	Val	Glu	Met	Leu	Pro	Ser
1760						1765					1770			

Lys	Val	Gly	Ile	Trp	Arg	Ile	Glu	Cys	Leu	Ile	Gly	Glu	His	Leu
1775						1780					1785			
Gln	Ala	Gly	Met	Ser	Thr	Thr	Phe	Leu	Val	Tyr	Ser	Lys	Glu	Cys
1790						1795					1800			
Gln	Ala	Pro	Leu	Gly	Met	Ala	Ser	Gly	Arg	Ile	Arg	Asp	Phe	Gln
1805						1810					1815			
Ile	Thr	Ala	Ser	Gly	Gln	Tyr	Gly	Gln	Trp	Ala	Pro	Lys	Leu	Ala
1820						1825					1830			
Arg	Leu	His	Tyr	Ser	Gly	Ser	Ile	Asn	Ala	Trp	Ser	Thr	Lys	Asp
1835						1840					1845			
Pro	His	Ser	Trp	Ile	Lys	Val	Asp	Leu	Leu	Ala	Pro	Met	Ile	Ile
1850						1855					1860			
His	Gly	Ile	Met	Thr	Gln	Gly	Ala	Arg	Gln	Lys	Phe	Ser	Ser	Leu
1865						1870					1875			
Tyr	Ile	Ser	Gln	Phe	Ile	Ile	Met	Tyr	Ser	Leu	Asp	Gly	Arg	Asn
1880						1885					1890			
Trp	Gln	Ser	Tyr	Arg	Gly	Asn	Ser	Thr	Gly	Thr	Leu	Met	Val	Phe
1895						1900					1905			
Phe	Gly	Asn	Val	Asp	Ala	Ser	Gly	Ile	Lys	His	Asn	Ile	Phe	Asn
1910						1915					1920			
Pro	Pro	Ile	Val	Ala	Arg	Tyr	Ile	Arg	Leu	His	Pro	Thr	His	Tyr
1925						1930					1935			
Ser	Ile	Arg	Ser	Thr	Leu	Arg	Met	Glu	Leu	Met	Gly	Cys	Asp	Leu
1940						1945					1950			
Asn	Ser	Cys	Ser	Met	Pro	Leu	Gly	Met	Gln	Asn	Lys	Ala	Ile	Ser
1955						1960					1965			
Asp	Ser	Gln	Ile	Thr	Ala	Ser	Ser	His	Leu	Ser	Asn	Ile	Phe	Ala
1970						1975					1980			
Thr	Trp	Ser	Pro	Ser	Gln	Ala	Arg	Leu	His	Leu	Gln	Gly	Arg	Thr
1985						1990					1995			
Asn	Ala	Trp	Arg	Pro	Arg	Val	Ser	Ser	Ala	Glu	Glu	Trp	Leu	Gln

2000	2005	2010
Val Asp Leu Gln Lys Thr 2015	Val Lys Val Thr Gly 2020	Ile Thr Thr Gln 2025
Gly Val Lys Ser Leu Leu 2030	Ser ¹ Met Tyr Val 2035	Lys Glu Phe Leu 2040
Val Ser Ser Ser Gln Asp 2045	Gly Arg Arg Trp Thr 2050	Leu Phe Leu Gln 2055
Asp Gly His Thr Lys Val 2060	Phe Gln Gly Asn Gln 2065	Asp Ser Ser Thr 2070
Pro Val Val Asn Ala Leu 2075	Asp Pro Pro Leu Phe 2080	Thr Arg Tyr Leu 2085
Arg Ile His Pro Thr Ser 2090	Trp Ala Gln His Ile 2095	Ala Leu Arg Leu 2100
Glu Val Leu Gly Cys Glu 2105	Ala Gln Asp Leu Tyr 2110	

<210> 4

<211> 6539

<212> DNA

<213> Sus scrofa

<400> 4

```

ctaaaatcct ttgtgaaaaa tttggggcctt ttcacgaaat cagaaaaagt ttaccttttc      60
tcccgggagc caaaaatagc ttagagaaga attcatcttt catttctcca gctggacatt      120
tttgatcagc aagagtcagc cagctagagc tctccacctg tgtctttctg tgtctcttgc      180
cactcggcct tagtgccatc aggagatact acctgggcgc agtggaactg tcctgggact      240
accggcaaag tgaactcctc cgtgagctgc acgtggacac cagatttcct gctacagcgc      300
caggagctct tccgttgggc ccgtcagtc tgtacaaaaa gactgtgttc gtagagttca      360
cggatcaact tttcagcgtt gccaggccca ggccaccatg gatgggtctg ctgggtccta      420
ccatccaggc tgagggtttac gacacgggtg tcgttaccct gaagaacatg gcttctcatc      480
ccgttagtct tcacgctgtc ggcgtctcct tctggaaatc ttccgaaggc gctgaatatg      540
aggatcacac cagccaaagg gagaaggaag acgataaagt ccttcccggc aaaagccaaa      600
cctacgtctg gcaggtcctg aaagaaaatg gtccaacagc ctctgaccca ccatgtctca      660
cctactcata cctgtctcac gtggacctgg tgaaagacct gaattcgggc ctcatggag      720

```

ccctgctggt ttgtagagaa gggagtctga ccagagaaag gacccagaac ctgcacgaat	780
ttgtactact ttttgctgtc tttgatgaag ggaaaagttg gcactcagca agaaatgact	840
cctggacacg ggccatggat cccgcacctg ccaggggccca gcctgcaatg cacacagtca	900
atggctatgt caacaggtct ctgccaggtc tgatcggatg tcataagaaa tcagtctact	960
ggcacgtgat tggaatgggc accagcccgg aagtgcactc catttttctt gaaggccaca	1020
cgtttctcgt gaggcacat cggcaggctt ccttgagat ctgccacta actttcctca	1080
ctgctcagac attcctgatg gaccttggcc agttcctact gttttgtcat atctcttccc	1140
accaccatgg tggcatggag gctcacgtca gagtagaaag ctgcgccgag gagccccagc	1200
tgcggaggaa agctgatgaa gaggaagatt atgatgacaa tttgtacgac tcggacatgg	1260
acgtggtccg gctcgatggg gacgacgtgt ctccctttat ccaaaccgc tcggttgcca	1320
agaagcatcc caaaacctgg gtgcactaca tctctgcaga ggaggaggac tgggactacg	1380
ccccgcggg cccagcccc agtgacagaa gttataaaaag tctctacttg aacagtgggc	1440
ctcagcgaat tggtaggaaa tacaaaaaag ctcgattcgt cgcttacacg gatgtaacat	1500
ttaagactcg taaagctatt ccgtatgaat caggaatcct gggaccttta ctttatggag	1560
aagttggaga cacacttttg attatattta agaataaagc gagccgacca tataacatct	1620
accctcatgg aatcactgat gtcagcgctt tgcaccagc gagacttcta aaaggttga	1680
aacatttgaa agacatgcca attctgccag gagagacttt caagtataaa tggacagtga	1740
ctgtggaaga tgggccaacc aagtccgac ctccggtgcct gaccgcctac tactcgagct	1800
ccattaatct agagaaagat ctggcttcgg gactcattgg ccctctcctc atctgctaca	1860
aagaatctgt agaccaaaga ggaaaccaga tgatgtcaga caagagaaac gtcacctgt	1920
tttctgtatt cgatgagaat caaagctggg acctgcaga gaatattcag cgcttcctcc	1980
ccaatccgga tggattacag cccaggatc cagagttcca agcttctaac atcatgcaca	2040
gcatcaatgg ctatgttttt gatagcttgc agctgtcggg ttgtttgcac gaggtggcat	2100
actggtacat tctaagtgtt ggagcacaga cggacttcct ctccgtcttc ttctctggct	2160
acaccttcaa acacaaaatg gtctatgaag acacactcac cctgttcccc ttctcaggag	2220
aaacggtcct catgtcaatg gaaaaccag gtctctgggt cctaggggtgc cacaactcag	2280
acttgccgaa cagagggatg acagccttac tgaaggtgta tagttgtgac agggacattg	2340
gtgattatta tgacaacact tatgaagata ttccaggctt cttgctgagt ggaaagaatg	2400
tcattgaacc cagaagcttt gccagaatt caagaccccc tagtgcgagc caaaagcaat	2460
tccaaaccat cacaagtcca gaagatgacg tggagcttga cccgcagtct ggagagagaa	2520
cccaagcact ggaagaacta agtgtccct ctggatgatg gtcgatgctc ttgggacaga	2580

atcctgctcc	acatgggtca	tcctcatctg	atcttcaaga	agccaggaat	gaggctgatg	2640
attatttacc	tggagcaaga	gaaagaaaca	cggcccatc	cgcagcggca	cgtctcagac	2700
cagagctgca	tcacagtgcc	gaaagagtac	ttactcctga	gccagagaaa	gagttgaaga	2760
aacttgattc	aaaaatgtct	agttcatcag	accttctaaa	gacttcgcca	acaattccat	2820
cagacacggt	gtcagcggag	actgaaagga	cacattcctt	aggcccccca	caccgcgagg	2880
ttaatttcag	gagtcaatta	ggtgccattg	tacttggcaa	aaattcatct	cactttattg	2940
gggctgggtg	ccctttgggc	tcgactgagg	aggatcatga	aagctccctg	ggagaaaatg	3000
tatcaccagt	ggagagtgc	gggatatttg	aaaaggaaa	agctcatgga	cctgcttcac	3060
tgaccaaaaga	cgatgtttta	tttaaagtta	atatctcttt	ggtaaagaca	aacaaggcac	3120
gagtttactt	aaaaactaat	agaaagattc	acattgatga	cgcagcttta	ttaactgaga	3180
atagggcatc	tgcaacgttt	atggacaaaa	atactacagc	ttcgggatta	aatcatgtgt	3240
caaattggat	aaaagggccc	cttggaaga	acccctaag	ctcggagcga	ggccccagtc	3300
cagagcttct	gacatcttca	ggatcaggaa	aatctgtgaa	aggtcagagt	tctgggcagg	3360
ggagaatacg	ggtggcagtg	gaagaggaag	aactgagcaa	aggcaaagag	atgatgcttc	3420
ccaacagcga	gtcaccttt	ctcactaact	cggctgatgt	ccaaggaaac	gatacacaca	3480
gtcaaggaaa	aaagtctcgg	gaagagatgg	aaaggagaga	aaaattagtc	caagaaaaag	3540
tcgacttgcc	tcaggtgtat	acagcgactg	gaactaagaa	tttcctgaga	aacatttttc	3600
accaaagcac	tgagcccagt	gtagaagggt	ttgatggggg	gtcacatgcg	ccggtgcctc	3660
aagacagcag	gtcattaaat	gattcggcag	agagagcaga	gactcacata	gcccatttct	3720
cagcaattag	ggaagaggca	cccttggaag	ccccgggaaa	tcgaacaggt	ccaggtccga	3780
ggagtgcggt	ccccgcgcg	gttaagcaga	gcttgaaaca	gatcagactc	ccgctagaag	3840
aaataaagcc	tgaagggggg	gtggttctga	atgccacctc	aaccgggtgg	tctgaaagca	3900
gtcctatctt	acaaggagcc	aaaagaaata	acctttcttt	acctttcctg	accttggaag	3960
tggccggagg	tcaaggaaag	atcagcgccc	tggggaaaag	tgccgcaggc	ccgctggcgt	4020
ccgggaagct	ggagaaggct	gttctctctt	cagcaggctt	gtctgaagca	tctggcaaag	4080
ctgagtttct	tcctaaagtt	cgagttcatc	gggaagacct	gttgctcaa	aaaaccagca	4140
atgtttcttg	cgcacacggg	gatctcggcc	aggagatctt	cctgcagaaa	acacggggac	4200
ctgttaacct	gaacaaagta	aatagacctg	gaaggactcc	ctccaagctt	ctgggtcccc	4260
cgatgcccaa	agagtgggaa	tccttagaga	agtcaccaa	aagcacagct	ctcaggacga	4320
aagacatcat	cagtttacct	ctggaccgtc	acgaaagcaa	tcattcaata	gcagcaaaaa	4380

atgaaggaca agccgagacc caaagagaag ccgcctggac gaagcaggga gggcctggaa	4440
ggctgtgcgc tccaaagcct ccggtcctgc gacggcatca gaggacata agccttccta	4500
cttttcagcc ggaggaagac aaaatggact atgatgatat cttctcaact gaaacgaagg	4560
gagaagatth tgacatthac ggtgaggatg aaaatcagga ccctcgcagc tttcagaaga	4620
gaacccgaca ctatthcatt gctgcggtgg agcagctctg ggattacggg atgagcgaat	4680
ccccccgggc gctaagaaac agggctcaga acggagaggt gcctcggttc aagaagggtg	4740
tcttccggga atthgctgac ggctccttca cgcagccgtc gtaccgctgg gaactcaaca	4800
aacacttggg gctcttggga ccctacatca gagcgggaagt tgaagacaac atcatggtaa	4860
ctttcaaaaa ccaggcgtct cgtccctatt ccttctactc gagccttatt tcttatccgg	4920
atgatcagga gcaaggggca gaacctcgac acaacttcgt ccagccaaat gaaaccagaa	4980
cttactthtg gaaagtgcag catcacatgg caccacaga agacgagtht gactgcaaag	5040
cctgggccta cttttctgat gttgacctgg aaaaagatgt gcactcaggc ttgatcggcc	5100
cccttctgat ctgccgcgcc aacacctga acgctgctca cggtagacaa gtgacctgc	5160
aagaatthtg tctgtthttc actatthttg atgagacaaa gagctggtac ttcactgaaa	5220
atgtggaaag gaactgccgg gccccctgcc acctgcagat ggaggacccc actctgaaag	5280
aaaactatcg cttccatgca atcaatggct atgtgatgga tacactccct ggcttagtaa	5340
tggctcagaa tcaaaggatc cgatggtatc tgctcagcat gggcagcaat gaaaatatcc	5400
attcgattca tthtagcgga cactgtttca gtgtacggaa aaaggaggag tataaaatgg	5460
ccgtgtacaa tctctatccg ggtgtctthg agacagtgga aatgctaccg tccaaagthg	5520
gaatthggcg aatagaatgc ctgattggcg agcacctgca agctgggatg agcacgactt	5580
tcctggtgta cagcaaggag tgtcaggctc cactgggaat ggcttctgga cgcattagag	5640
atthtcagat cacagcttca ggacagtatg gacagtgggc cccaaagctg gccagacttc	5700
attatthccg atcaatcaat gcctggagca ccaaggatcc cactcctgg atcaaggthg	5760
atctgthggc accaatgatc attcacggca tcatgaccca gggtgcccgt cagaagthtt	5820
ccagcctcta catctcccag thtatcatca tgtacagtct tgacgggagg aactggcaga	5880
gttaccgagg gaatthcacg ggcaccttaa tggctcttct tggcaatgtg gacgcatctg	5940
ggattaaaca caatatthtt aacctccga thgtggctcg gtacatccgt thgcacccaa	6000
cacattacag catccgcagc actcttcgca tggagthgat gggctgtgat ttaaacagth	6060
gcagcatgcc cctgggaatg cagaataaag cgatatcaga ctcacagatc acggcctcct	6120
cccacctaag caatatatth gccacctggc ctcttcaca agcccgactt cacctccagg	6180
ggcggacgaa tgcttgccga ccccggtga gcagcgcaga ggagthggctg caggtggacc	6240

tgcagaagac	ggtgaaggtc	acaggcatca	ccaccacagg	cgtgaagtcc	ctgctcagca	6300
gcatgtatgt	gaaggagttc	ctcgtgtcca	gtagtcagga	cggccgccgc	tggaccctgt	6360
ttcttcagga	cggccacacg	aaggtttttc	agggcaatca	ggactcctcc	acccccgtgg	6420
tgaacgctct	ggaccccccg	ctgttcacgc	gctacctgag	gatccacccc	acgagctggg	6480
cgcagcacat	cgccctgagg	ctcgagggtc	taggatgtga	ggcacaggat	ctctactga	6539